

# Contents

|                                                                                                       |          |                                                                                                                                                               |           |
|-------------------------------------------------------------------------------------------------------|----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------|
| <b>Chapter 1 Cells, Genomes, and the Diversity of Life</b>                                            | <b>1</b> |                                                                                                                                                               |           |
| <b>THE UNIVERSAL FEATURES OF LIFE ON EARTH</b>                                                        |          |                                                                                                                                                               |           |
| All Cells Store Their Hereditary Information in the Form of Double-Strand DNA Molecules               | 2        | The Expression Levels of All the Genes of an Organism Can Be Determined                                                                                       | 37        |
| All Cells Replicate Their Hereditary Information by Templated Polymerization                          | 2        | <i>Arabidopsis</i> Has Been Chosen as a Model Plant                                                                                                           | 38        |
| All Cells Transcribe Portions of Their DNA into RNA Molecules                                         | 3        | The World of Animal Cells Is Mainly Represented by a Worm, a Fly, a Fish, a Mouse, and a Human                                                                | 38        |
| All Cells Use Proteins as Catalysts                                                                   | 5        | Studies in the Fruit Fly <i>Drosophila</i> Provide a Key to Vertebrate Development                                                                            | 39        |
| All Cells Translate RNA into Protein in the Same Way                                                  | 6        | The Frog and the Zebrafish Provide Highly Accessible Vertebrate Models                                                                                        | 40        |
| Each Protein Is Encoded by a Specific Gene                                                            | 6        | The Mouse Is the Predominant Mammalian Model Organism                                                                                                         | 41        |
| Life Requires a Continual Input of Free Energy                                                        | 7        | The COVID-19 Pandemic Has Focused Scientists on the SARS-CoV-2 Coronavirus                                                                                    | 42        |
| All Cells Function as Biochemical Factories                                                           | 7        | Humans Are Unique in Reporting on Their Own Peculiarities To Understand Cells and Organisms Will Require Mathematics, Computers, and Quantitative Information | 44        |
| All Cells Are Enclosed in a Plasma Membrane Across Which Nutrients and Waste Materials Must Pass      | 8        | Summary                                                                                                                                                       | 45        |
| Cells Operate at a Microscopic Scale Dominated by Random Thermal Motion                               | 9        | Problems                                                                                                                                                      | 46        |
| A Living Cell Can Exist with 500 Genes                                                                | 10       | References                                                                                                                                                    | 47        |
| Summary                                                                                               | 10       |                                                                                                                                                               |           |
| <b>GENOME DIVERSIFICATION AND THE TREE OF LIFE</b>                                                    |          | <b>Chapter 2 Cell Chemistry and Bioenergetics</b>                                                                                                             | <b>49</b> |
| The Tree of Life Has Three Major Domains: Eukaryotes, Bacteria, and Archaea                           | 11       | <b>THE CHEMICAL COMPONENTS OF A CELL</b>                                                                                                                      | <b>49</b> |
| Eukaryotes Make Up the Domain of Life That Is Most Familiar to Us                                     | 13       | Water Is Held Together by Hydrogen Bonds                                                                                                                      | 50        |
| On the Basis of Genome Analysis, Bacteria Are the Most Diverse Group of Organisms on the Planet       | 13       | Four Types of Noncovalent Attractions Help Bring Molecules Together in Cells                                                                                  | 51        |
| Archaea: The Most Mysterious Domain of Life                                                           | 15       | Some Polar Molecules Form Acids and Bases in Water                                                                                                            | 52        |
| Organisms Occupy Most of Our Planet                                                                   | 15       | A Cell Is Formed from Carbon Compounds                                                                                                                        | 53        |
| Cells Can Be Powered by a Wide Variety of Free-Energy Sources                                         | 15       | Cells Contain Four Major Families of Small Organic Molecules                                                                                                  | 53        |
| Some Cells Fix Nitrogen and Carbon Dioxide for Other Cells                                            | 17       | The Chemistry of Cells Is Dominated by Macromolecules with Remarkable Properties                                                                              | 54        |
| Genomes Diversify Over Evolutionary Time, Producing New Types of Organisms                            | 18       | Noncovalent Bonds Specify Both the Precise Shape of a Macromolecule and Its Binding to Other Molecules                                                        | 55        |
| New Genes Are Generated from Preexisting Genes                                                        | 19       | Summary                                                                                                                                                       | 56        |
| Gene Duplications Give Rise to Families of Related Genes Within a Single Genome                       | 20       | <b>CATALYSIS AND THE USE OF ENERGY BY CELLS</b>                                                                                                               | <b>57</b> |
| The Function of a Gene Can Often Be Deduced from Its Nucleotide Sequence                              | 20       | Cell Metabolism Is Organized by Enzymes                                                                                                                       | 57        |
| More Than 200 Gene Families Are Common to All Three Domains of Life                                   | 21       | Biological Order Is Made Possible by the Release of Heat Energy from Cells                                                                                    | 58        |
| Summary                                                                                               | 21       | Cells Obtain Energy by the Oxidation of Organic Molecules                                                                                                     | 61        |
| <b>EUKARYOTES AND THE ORIGIN OF THE EUKARYOTIC CELL</b>                                               |          | Oxidation and Reduction Involve Electron Transfers                                                                                                            | 62        |
| Eukaryotic Cells Contain a Variety of Organelles                                                      | 22       | Enzymes Lower the Activation-Energy Barriers That Block Chemical Reactions                                                                                    | 63        |
| Mitochondria Evolved from a Symbiotic Bacterium Captured by an Ancient Archaeon                       | 23       | Enzymes Can Drive Substrate Molecules Along Specific Reaction Pathways                                                                                        | 64        |
| Chloroplasts Evolved from a Symbiotic Photosynthetic Bacterium Engulfed by an Ancient Eukaryotic Cell | 25       | How Enzymes Find Their Substrates: The Enormous Rapidity of Molecular Motions                                                                                 | 65        |
| Eukaryotes Have Hybrid Genomes                                                                        | 26       | The Free-Energy Change for a Reaction, $\Delta G$ , Determines Whether It Can Occur Spontaneously                                                             | 66        |
| Eukaryotic Genomes Are Big                                                                            | 27       | The Concentration of Reactants Influences the Free-Energy Change and a Reaction's Direction                                                                   | 67        |
| Eukaryotic Genomes Are Rich in Regulatory DNA                                                         | 28       | The Standard Free-Energy Change, $\Delta G^\circ$ , Makes It Possible to Compare the Energetics of Different Reactions                                        | 67        |
| Eukaryotic Genomes Define the Program of Multicellular Development                                    | 29       | The Equilibrium Constant and $\Delta G^\circ$ Are Readily Derived from Each Other                                                                             | 68        |
| Many Eukaryotes Live as Solitary Cells                                                                | 30       | The Free-Energy Changes of Coupled Reactions Are Additive                                                                                                     | 69        |
| Summary                                                                                               | 31       | Activated Carrier Molecules Are Essential for Biosynthesis                                                                                                    | 69        |
| <b>MODEL ORGANISMS</b>                                                                                |          | The Formation of an Activated Carrier Is Coupled to an Energetically Favorable Reaction                                                                       | 70        |
| Mutations Reveal the Functions of Genes                                                               | 31       | ATP Is the Most Widely Used Activated Carrier Molecule                                                                                                        | 71        |
| Molecular Biology Began with a Spotlight on One Bacterium and Its Viruses                             | 32       | Energy Stored in ATP Is Often Harnessed to Join Two Molecules Together                                                                                        | 72        |
| The Focus on <i>E. coli</i> as a Model Organism Has Accelerated Many Subsequent Discoveries           | 33       |                                                                                                                                                               |           |
| A Yeast Serves as a Minimal Model Eukaryote                                                           | 35       |                                                                                                                                                               |           |
|                                                                                                       | 36       |                                                                                                                                                               |           |

|                                                                    |     |                                                                  |     |
|--------------------------------------------------------------------|-----|------------------------------------------------------------------|-----|
| NADH and NADPH Are Important Electron Carriers                     | 73  | Enzymes Speed Reactions by Selectively Stabilizing               |     |
| There Are Many Other Activated Carrier Molecules in Cells          | 75  | Transition States                                                | 148 |
| The Synthesis of Biological Polymers Is Driven by                  |     | Enzymes Can Use Simultaneous Acid and Base Catalysis             | 148 |
| ATP Hydrolysis                                                     | 76  | Lysozyme Illustrates How an Enzyme Works                         | 149 |
| Summary                                                            | 78  | Tightly Bound Small Molecules Add Extra Functions to Proteins    | 152 |
| <b>HOW CELLS OBTAIN ENERGY FROM FOOD</b>                           | 80  | The Cell Regulates the Catalytic Activities of Its Enzymes       | 155 |
| Glycolysis Is a Central ATP-producing Pathway                      | 80  | Allosteric Enzymes Have Two or More Binding Sites That Interact  | 155 |
| Glycolysis Illustrates How Enzymes Couple Oxidation                |     | Two Ligands Whose Binding Sites Are Coupled Must Reciprocally    |     |
| to Energy Storage                                                  | 83  | Affect Each Other's Binding                                      | 157 |
| Fermentations Produce ATP in the Absence of Oxygen                 | 84  | Symmetrical Protein Assemblies Produce Cooperative Allosteric    |     |
| Organisms Store Food Molecules in Special Reservoirs               | 85  | Transitions                                                      | 158 |
| Between Meals, Most Animal Cells Derive Their Energy               |     | Many Changes in Proteins Are Driven by Protein Phosphorylation   | 159 |
| from Fatty Acids Obtained from Fat                                 | 86  | A Eukaryotic Cell Contains a Large Collection of Protein Kinases |     |
| Sugars and Fats Are Both Degraded to Acetyl CoA                    |     | and Protein Phosphatases                                         | 159 |
| in Mitochondria                                                    | 87  | The Regulation of the Src Protein Kinase Reveals How a           |     |
| The Citric Acid Cycle Generates NADH by Oxidizing                  |     | Protein Can Function as a Microprocessor                         | 161 |
| Acetyl Groups to CO <sub>2</sub>                                   | 88  | Regulatory GTP-binding Proteins Are Switched On and              |     |
| Electron Transport Drives the Synthesis of the Majority            |     | Off by the Gain and Loss of a Phosphate Group                    | 162 |
| of the ATP in Most Cells                                           | 90  | Proteins Can Be Regulated by the Covalent Addition               |     |
| Many Biosynthetic Pathways Begin with Glycolysis                   |     | of Other Proteins                                                | 162 |
| or the Citric Acid Cycle                                           | 90  | An Elaborate Ubiquitin-conjugating System Is Used                |     |
| Animals Must Obtain All the Nitrogen and Sulfur                    |     | to Mark Proteins                                                 | 163 |
| They Need from Food                                                | 91  | Protein Complexes with Interchangeable Parts Make                |     |
| Metabolism Is Highly Organized and Regulated                       | 92  | Efficient Use of Genetic Information                             | 164 |
| Summary                                                            | 93  | A GTP-binding Protein Shows How Large Protein                    |     |
| Problems                                                           | 112 | Movements Can Be Generated from Small Ones                       | 166 |
| References                                                         | 114 | Motor Proteins Produce Directional Movement in Cells             | 167 |
|                                                                    |     | Proteins Often Form Large Complexes That Function as Protein     |     |
|                                                                    |     | Machines                                                         | 167 |
| <b>Chapter 3 Proteins</b>                                          | 115 | The Disordered Regions in Proteins Are Critical for a            |     |
|                                                                    |     | Set of Different Functions                                       | 168 |
| <b>THE ATOMIC STRUCTURE OF PROTEINS</b>                            | 115 | Scaffolds Bring Sets of Interacting Macromolecules Together      |     |
| The Structure of a Protein Is Specified by Its Amino               |     | and Concentrate Them in Selected Regions of a Cell               | 170 |
| Acid Sequence                                                      | 115 | Macromolecules Can Self-assemble to Form Biomolecular            |     |
| Proteins Fold into a Conformation of Lowest Energy                 | 121 | Condensates                                                      | 171 |
| The $\alpha$ Helix and the $\beta$ Sheet Are Common Folding Motifs | 121 | Classical Studies of Phase Separation Have Relevance             |     |
| Four Levels of Organization Are Considered to Contribute           |     | for Biomolecular Condensates                                     | 173 |
| to Protein Structure                                               | 123 | A Comparison of Three Important Types of Large Biological        |     |
| Protein Domains Are the Modular Units from Which Larger            |     | Assemblies                                                       | 174 |
| Proteins Are Built                                                 | 124 | Many Proteins Are Controlled by Covalent Modifications           |     |
| Proteins Must Contain Unstructured Regions                         | 126 | That Direct Them to Specific Sites Inside the Cell               | 175 |
| All Protein Structures Are Dynamic, Interconverting Rapidly        |     | A Complex Network of Protein Interactions Underlies              |     |
| Between an Ensemble of Closely Related Conformations               |     | Cell Function                                                    | 176 |
| Because of Thermal Energy                                          | 126 | Protein Structures Can Be Predicted and New Proteins             |     |
| Function Has Selected for a Tiny Fraction of the Many              |     | Designed                                                         | 178 |
| Possible Polypeptide Chains                                        | 126 | Summary                                                          | 179 |
| Proteins Can Be Classified into Many Families                      | 127 | Problems                                                         | 179 |
| Some Protein Domains Are Found in Many Different Proteins          | 129 | References                                                       | 181 |
| The Human Genome Encodes a Complex Set of Proteins,                |     |                                                                  |     |
| Revealing That Much Remains Unknown                                | 130 |                                                                  |     |
| Protein Molecules Often Contain More Than One                      |     |                                                                  |     |
| Polypeptide Chain                                                  | 130 |                                                                  |     |
| Some Globular Proteins Form Long Helical Filaments                 | 131 |                                                                  |     |
| Protein Molecules Can Have Elongated, Fibrous Shapes               | 132 |                                                                  |     |
| Covalent Cross-Linkages Stabilize Extracellular Proteins           | 133 |                                                                  |     |
| Protein Molecules Often Serve as Subunits for the Assembly         |     |                                                                  |     |
| of Large Structures                                                | 134 |                                                                  |     |
| Many Structures in Cells Are Capable of Self-Assembly              | 136 |                                                                  |     |
| Assembly Factors Often Aid the Formation of Complex                |     |                                                                  |     |
| Biological Structures                                              | 136 |                                                                  |     |
| When Assembly Processes Go Wrong: The Case of                      |     |                                                                  |     |
| Amyloid Fibrils                                                    | 137 |                                                                  |     |
| Amyloid Structures Can Also Perform Useful Functions in Cells      | 139 |                                                                  |     |
| Summary                                                            | 140 |                                                                  |     |
| <b>PROTEIN FUNCTION</b>                                            | 140 |                                                                  |     |
| All Proteins Bind to Other Molecules                               | 140 |                                                                  |     |
| The Surface Conformation of a Protein Determines Its Chemistry     | 142 |                                                                  |     |
| Sequence Comparisons Between Protein Family Members                |     |                                                                  |     |
| Highlight Crucial Ligand-binding Sites                             | 142 |                                                                  |     |
| Proteins Bind to Other Proteins Through Several Types              |     |                                                                  |     |
| of Interfaces                                                      | 143 |                                                                  |     |
| Antibody Binding Sites Are Especially Versatile                    | 144 |                                                                  |     |
| The Equilibrium Constant Measures Binding Strength                 | 145 |                                                                  |     |
| Enzymes Are Powerful and Highly Specific Catalysts                 | 146 |                                                                  |     |
| Substrate Binding Is the First Step in Enzyme Catalysis            | 146 |                                                                  |     |
|                                                                    |     | <b>Chapter 4 DNA, Chromosomes, and Genomes</b>                   | 183 |
|                                                                    |     | <b>THE STRUCTURE AND FUNCTION OF DNA</b>                         | 185 |
|                                                                    |     | A DNA Molecule Consists of Two Complementary Chains of           |     |
|                                                                    |     | Nucleotides                                                      | 185 |
|                                                                    |     | The Structure of DNA Provides a Mechanism for Heredity           | 187 |
|                                                                    |     | In Eukaryotes, DNA Is Enclosed in a Cell Nucleus                 | 189 |
|                                                                    |     | Summary                                                          | 189 |
|                                                                    |     | <b>CHROMOSOMAL DNA AND ITS PACKAGING</b>                         |     |
|                                                                    |     | <b>IN THE CHROMATIN FIBER</b>                                    | 189 |
|                                                                    |     | Eukaryotic DNA Is Packaged into a Set of Chromosomes             | 190 |
|                                                                    |     | Chromosomes Contain Long Strings of Genes                        | 191 |
|                                                                    |     | The Nucleotide Sequence of the Human Genome Shows                |     |
|                                                                    |     | How Our Genes Are Arranged                                       | 193 |
|                                                                    |     | Each DNA Molecule That Forms a Linear Chromosome                 |     |
|                                                                    |     | Must Contain a Centromere, Two Telomeres,                        |     |
|                                                                    |     | and Replication Origins                                          | 195 |
|                                                                    |     | DNA Molecules Are Highly Condensed in Chromosomes                | 197 |
|                                                                    |     | Nucleosomes Are a Basic Unit of Eukaryotic Chromosome            |     |
|                                                                    |     | Structure                                                        | 197 |
|                                                                    |     | The Structure of the Nucleosome Core Particle Reveals            |     |
|                                                                    |     | How DNA Is Packaged                                              | 198 |
|                                                                    |     | Nucleosomes Have a Dynamic Structure and Are                     |     |
|                                                                    |     | Frequently Subjected to Changes Catalyzed by                     |     |
|                                                                    |     | ATP-dependent Chromatin-remodeling Complexes                     | 200 |

|                                                                                                                                                 |     |                                                                                                                        |            |
|-------------------------------------------------------------------------------------------------------------------------------------------------|-----|------------------------------------------------------------------------------------------------------------------------|------------|
| Attractions Between Nucleosomes Compact the Chromatin Fiber                                                                                     | 202 | We Can Trace Human History by Analyzing Genomes                                                                        | 244        |
| Summary                                                                                                                                         | 203 | The Sequencing of Hundreds of Thousands of Human Genomes Reveals Much Variation                                        | 245        |
| <b>THE EFFECT OF CHROMATIN STRUCTURE ON DNA FUNCTION</b>                                                                                        | 203 | Most of the Variants Observed in the Human Population Are Common Alleles, with at Most a Weak Effect on Phenotype      | 246        |
| Different Regions of the Human Genome Are Packaged Very Differently in Chromatin                                                                | 204 | Forensic Analyses Exploit Special DNA Sequences with Unusually High Mutation Rates                                     | 247        |
| Heterochromatin Is Highly Condensed and Restricts Gene Expression                                                                               | 204 | An Understanding of Human Variation Is Critical for Improving Medicine                                                 | 248        |
| The Heterochromatic State Can Spread Along a Chromosome and Be Inherited from One Cell Generation to the Next                                   | 205 | Summary                                                                                                                | 248        |
| The Core Histones Are Covalently Modified at Many Different Sites                                                                               | 206 | Problems                                                                                                               | 249        |
| Chromatin Acquires Additional Variety Through the Site-specific Insertion of a Small Set of Histone Variants                                    | 208 | References                                                                                                             | 251        |
| Covalent Modifications and Histone Variants Can Act in Concert to Control Chromosome Functions                                                  | 208 | <b>Chapter 5 DNA Replication, Repair, and Recombination</b>                                                            | <b>253</b> |
| A Complex of Reader and Writer Proteins Can Spread Specific Chromatin Modifications Along a Chromosome                                          | 210 | <b>THE MAINTENANCE OF DNA SEQUENCES</b>                                                                                | <b>253</b> |
| Barrier DNA-Protein Complexes Block the Spread of Reader-Writer Complexes and Thereby Separate Neighboring Chromatin Domains                    | 212 | Mutation Rates Are Extremely Low                                                                                       | 253        |
| Centromeres Have a Special, Inherited Chromatin Structure                                                                                       | 213 | Low Mutation Rates Are Necessary for Life as We Know It                                                                | 254        |
| Some Forms of Chromatin Can Be Directly Inherited                                                                                               | 215 | Summary                                                                                                                | 255        |
| The Abnormal Perturbations of Heterochromatin That Arise During Tumor Progression Contribute to Many Cancers                                    | 215 | <b>DNA REPLICATION MECHANISMS</b>                                                                                      | <b>255</b> |
| Summary                                                                                                                                         | 217 | Base-pairing Underlies DNA Replication and DNA Repair                                                                  | 255        |
| <b>THE GLOBAL STRUCTURE OF CHROMOSOMES</b>                                                                                                      | 217 | The DNA Replication Fork Is Asymmetrical                                                                               | 256        |
| Chromosomes Are Folded into Large Loops of Chromatin                                                                                            | 217 | The High Fidelity of DNA Replication Requires Several Proofreading Mechanisms                                          | 258        |
| Polytene Chromosomes Are Uniquely Useful for Visualizing Chromatin Structures                                                                   | 218 | DNA Replication in the 5'-to-3' Direction Allows Efficient Error Correction                                            | 260        |
| Chromosome Loops Decondense When the Genes Within Them Are Expressed                                                                            | 220 | A Special Nucleotide-polymerizing Enzyme Synthesizes Short RNA Primer Molecules                                        | 260        |
| Mammalian Interphase Chromosomes Occupy Discrete Territories in the Nucleus, with Their Heterochromatin and Euchromatin Distributed Differently | 220 | Special Proteins Help to Open Up the DNA Double Helix in Front of the Replication Fork                                 | 261        |
| A Biochemical Technique Called Hi-C Reveals Details of Chromosome Organization                                                                  | 221 | A Sliding Ring Holds a Moving DNA Polymerase onto the DNA                                                              | 262        |
| Chromosomal DNA Is Organized into Loops by Large Protein Rings                                                                                  | 223 | The Proteins at a Replication Fork Cooperate to Form a Replication Machine                                             | 263        |
| Euchromatin and Heterochromatin Separate Spatially in the Nucleus                                                                               | 225 | DNA Replication Is Fundamentally Similar in Eukaryotes and Bacteria                                                    | 265        |
| Mitotic Chromosomes Are Highly Condensed                                                                                                        | 227 | A Strand-directed Mismatch Repair System Removes Replication Errors That Remain in the Wake of the Replication Machine | 267        |
| Summary                                                                                                                                         | 228 | The Accidental Incorporation of Ribonucleotides During DNA Replication Is Corrected                                    | 269        |
| <b>HOW GENOMES EVOLVE</b>                                                                                                                       | 229 | DNA Topoisomerases Prevent DNA Tangling During Replication                                                             | 269        |
| Genome Comparisons Reveal Functional DNA Sequences by Their Conservation Throughout Evolution                                                   | 230 | Summary                                                                                                                | 272        |
| Genome Alterations Are Caused by Failures of the Normal Mechanisms for Copying and Maintaining DNA, as Well as by Transposable DNA Elements     | 231 | <b>THE INITIATION AND COMPLETION OF DNA REPLICATION IN CHROMOSOMES</b>                                                 | <b>272</b> |
| The Genome Sequences of Two Species Differ in Proportion to the Length of Time Since They Have Separately Evolved                               | 232 | DNA Synthesis Begins at Replication Origins                                                                            | 272        |
| Phylogenetic Trees Constructed from a Comparison of DNA Sequences Trace the Relationships of All Organisms                                      | 233 | Bacterial Chromosomes Typically Have a Single Origin of DNA Replication                                                | 273        |
| A Comparison of Human and Mouse Chromosomes Shows How the Structures of Genomes Diverge                                                         | 234 | Eukaryotic Chromosomes Contain Multiple Origins of Replication                                                         | 273        |
| The Size of a Vertebrate Genome Reflects the Relative Rates of DNA Addition and DNA Loss in a Lineage                                           | 236 | In Eukaryotes, DNA Replication Takes Place During Only One Part of the Cell Cycle                                      | 276        |
| Multispecies Sequence Comparisons Identify Many Conserved DNA Sequences of Unknown Function                                                     | 237 | Eukaryotic Origins of Replication Are "Licensed" for Replication by the Assembly of an Origin Recognition Complex      | 276        |
| Changes in Previously Conserved Sequences Can Help Decipher Critical Steps in Evolution                                                         | 238 | Features of the Human Genome That Specify Origins of Replication Remain to Be Fully Understood                         | 277        |
| Mutations in the DNA Sequences That Control Gene Expression Have Driven Many of the Evolutionary Changes in Vertebrates                         | 239 | Properties of the ORC Ensure That Each Region of the DNA Is Replicated Once and Only Once in Each S Phase              | 277        |
| Gene Duplication Also Provides an Important Source of Genetic Novelty During Evolution                                                          | 240 | New Nucleosomes Are Assembled Behind the Replication Fork                                                              | 279        |
| Duplicated Genes Diverge                                                                                                                        | 240 | Termination of DNA Replication Occurs Through the Ordered Disassembly of the Replication Fork                          | 280        |
| The Evolution of the Globin Gene Family Shows How DNA Duplications Contribute to the Evolution of Organisms                                     | 241 | Telomerase Replicates the Ends of Chromosomes                                                                          | 281        |
| Genes Encoding New Proteins Can Be Created by the Recombination of Exons                                                                        | 242 | Telomeres Are Packaged into Specialized Structures That Protect the Ends of Chromosomes                                | 282        |
| Neutral Mutations Often Spread to Become Fixed in a Population, with a Probability That Depends on Population Size                              | 243 | Telomere Length Is Regulated by Cells and Organisms                                                                    | 282        |
|                                                                                                                                                 |     | Summary                                                                                                                | 284        |
|                                                                                                                                                 |     | <b>DNA REPAIR</b>                                                                                                      | <b>284</b> |
|                                                                                                                                                 |     | Without DNA Repair, Spontaneous DNA Damage Would Rapidly Change DNA Sequences                                          | 286        |
|                                                                                                                                                 |     | The DNA Double Helix Is Readily Repaired                                                                               | 288        |
|                                                                                                                                                 |     | DNA Damage Can Be Removed by More Than One Pathway                                                                     | 288        |

|                                                                                                                         |            |                                                                                                                                |            |
|-------------------------------------------------------------------------------------------------------------------------|------------|--------------------------------------------------------------------------------------------------------------------------------|------------|
| Coupling Nucleotide Excision Repair to Transcription Ensures That the Cell's Most Important DNA Is Efficiently Repaired | 290        | In Eukaryotes, Transcription Initiation Also Requires Activator, Mediator, and Chromatin-modifying Proteins                    | 334        |
| The Chemistry of the DNA Bases Facilitates Damage Detection                                                             | 290        | Transcription Elongation in Eukaryotes Requires Accessory Proteins                                                             | 335        |
| Special Translesion DNA Polymerases Are Used in Emergencies                                                             | 292        | Transcription Creates Superhelical Tension                                                                                     | 335        |
| Double-Strand Breaks Are Efficiently Repaired                                                                           | 292        | Transcription Elongation in Eukaryotes Is Tightly Coupled to RNA Processing                                                    | 337        |
| DNA Damage Delays Progression of the Cell Cycle                                                                         | 295        | RNA Capping Is the First Modification of Eukaryotic Pre-mRNAs                                                                  | 338        |
| Summary                                                                                                                 | 295        | RNA Splicing Removes Intron Sequences from Newly Transcribed Pre-mRNAs                                                         | 339        |
| <b>HOMOLOGOUS RECOMBINATION</b>                                                                                         | <b>296</b> | Nucleotide Sequences Signal Where Splicing Occurs                                                                              | 341        |
| Homologous Recombination Has Common Features in All Cells                                                               | 296        | RNA Splicing Is Performed by the Spliceosome                                                                                   | 341        |
| DNA Base-pairing Guides Homologous Recombination                                                                        | 296        | The Spliceosome Uses ATP Hydrolysis to Produce a Complex Series of RNA-RNA Rearrangements                                      | 343        |
| Homologous Recombination Can Flawlessly Repair Double-Strand Breaks in DNA                                              | 297        | Other Properties of Pre-mRNA and Its Synthesis Help to Explain the Choice of Proper Splice Sites                               | 345        |
| Specialized Processing of Double-Strand Breaks Commits Repair to Homologous Recombination                               | 298        | RNA Splicing Has Remarkable Plasticity                                                                                         | 346        |
| Strand Exchange Is Directed by the RecA/Rad51 Protein                                                                   | 298        | Spliceosome-catalyzed RNA Splicing Evolved from RNA Self-splicing Mechanisms                                                   | 347        |
| Homologous Recombination Can Rescue Broken and Stalled DNA Replication Forks                                            | 299        | RNA-processing Enzymes Generate the 3' End of Eukaryotic mRNAs                                                                 | 348        |
| DNA Repair by Homologous Recombination Entails Risks to the Cell                                                        | 300        | Mature Eukaryotic mRNAs Are Selectively Exported from the Nucleus                                                              | 349        |
| Homologous Recombination Is Crucial for Meiosis                                                                         | 301        | Noncoding RNAs Are Also Synthesized and Processed in the Nucleus                                                               | 351        |
| Meiotic Recombination Begins with a Programmed Double-Strand Break                                                      | 302        | The Nucleolus Is a Ribosome-producing Factory                                                                                  | 353        |
| Holliday Junctions Are Recognized by Enzymes That Drive Branch Migration                                                | 302        | The Nucleus Contains a Variety of Subnuclear Biomolecular Condensates                                                          | 355        |
| Homologous Recombination Produces Crossovers Between Maternal and Paternal Chromosomes During Meiosis                   | 304        | Summary                                                                                                                        | 357        |
| Homologous Recombination Often Results in Gene Conversion                                                               | 305        | <b>FROM RNA TO PROTEIN</b>                                                                                                     | <b>358</b> |
| Summary                                                                                                                 | 306        | An mRNA Sequence Is Decoded in Sets of Three Nucleotides                                                                       | 358        |
| <b>TRANSPPOSITION AND CONSERVATIVE SITE-SPECIFIC RECOMBINATION</b>                                                      | <b>306</b> | tRNA Molecules Match Amino Acids to Codons in mRNA                                                                             | 359        |
| Through Transposition, Mobile Genetic Elements Can Insert into Any DNA Sequence                                         | 307        | tRNAs Are Covalently Modified Before They Exit from the Nucleus                                                                | 361        |
| DNA-only Transposons Can Move by a Cut-and-Paste Mechanism                                                              | 307        | Specific Enzymes Couple Each Amino Acid to Its Appropriate tRNA Molecule                                                       | 361        |
| Some DNA-only Transposons Move by Replicating Themselves                                                                | 309        | Editing by tRNA Synthetases Ensures Accuracy                                                                                   | 363        |
| Some Viruses Use a Transposition Mechanism to Move Themselves into Host-Cell Chromosomes                                | 309        | Amino Acids Are Added to the C-terminal End of a Growing Polypeptide Chain                                                     | 364        |
| Some RNA Viruses Replicate and Express Their Genomes Without Using DNA as an Intermediate                               | 311        | The RNA Message Is Decoded in Ribosomes                                                                                        | 365        |
| Retroviral-like Retrotransposons Resemble Retroviruses, but Cannot Move from Cell to Cell                               | 313        | Elongation Factors Drive Translation Forward and Improve Its Accuracy                                                          | 368        |
| A Large Fraction of the Human Genome Is Composed of Nonretroviral Retrotransposons                                      | 313        | Induced Fit and Kinetic Proofreading Help Biological Processes Overcome the Inherent Limitations of Complementary Base-Pairing | 369        |
| Different Transposable Elements Predominate in Different Organisms                                                      | 314        | Accuracy in Translation Requires a Large Expenditure of Free Energy                                                            | 370        |
| Genome Sequences Reveal the Approximate Times at Which Transposable Elements Have Moved                                 | 314        | The Ribosome Is a Ribozyme                                                                                                     | 371        |
| Conservative Site-specific Recombination Can Reversibly Rearrange DNA                                                   | 315        | Nucleotide Sequences in mRNA Signal Where to Start Protein Synthesis                                                           | 373        |
| Conservative Site-specific Recombination Can Be Used to Turn Genes On or Off                                            | 316        | Stop Codons Mark the End of Translation                                                                                        | 374        |
| Bacterial Conservative Site-specific Recombinases Have Become Powerful Tools for Cell and Developmental Biologists      | 317        | Proteins Are Made on Polyribosomes                                                                                             | 375        |
| Summary                                                                                                                 | 317        | There Are Minor Variations in the Standard Genetic Code                                                                        | 375        |
| Problems                                                                                                                | 318        | Inhibitors of Prokaryotic Protein Synthesis Are Useful as Antibiotics                                                          | 376        |
| References                                                                                                              | 320        | Quality-Control Mechanisms Act to Prevent Translation of Damaged mRNAs                                                         | 378        |
|                                                                                                                         |            | Stalled Ribosomes Can Be Rescued                                                                                               | 379        |
|                                                                                                                         |            | The Ribosome Coordinates the Folding, Enzymatic Modification, and Assembly of Newly Synthesized Proteins                       | 380        |
| <b>Chapter 6 How Cells Read the Genome: From DNA to Protein</b>                                                         | <b>321</b> | Molecular Chaperones Help Guide the Folding of Most Proteins                                                                   | 380        |
| <b>FROM DNA TO RNA</b>                                                                                                  | <b>323</b> | Proper Folding of Newly Synthesized Proteins Is Also Aided by Translation Speed and Subunit Assembly                           | 383        |
| RNA Molecules Are Single-Stranded                                                                                       | 324        | Proteins That Ultimately Fail to Fold Correctly Are Marked for Destruction by Polyubiquitin                                    | 384        |
| Transcription Produces RNA Complementary to One Strand of DNA                                                           | 325        | The Proteasome Is a Compartmentalized Protease with Sequestered Active Sites                                                   | 384        |
| RNA Polymerases Carry Out DNA Transcription                                                                             | 325        | Many Proteins Are Controlled by Regulated Destruction                                                                          | 386        |
| Cells Produce Different Categories of RNA Molecules                                                                     | 327        | There Are Many Steps from DNA to Protein                                                                                       | 387        |
| Signals Encoded in DNA Tell RNA Polymerase Where to Start and Stop                                                      | 328        | Summary                                                                                                                        | 388        |
| Bacterial Transcription Start and Stop Signals Are Heterogeneous in Nucleotide Sequence                                 | 329        | <b>THE RNA WORLD AND THE ORIGINS OF LIFE</b>                                                                                   | <b>389</b> |
| Transcription Initiation in Eukaryotes Requires Many Proteins                                                           | 331        | Single-Strand RNA Molecules Can Fold into Highly Elaborate Structures                                                          | 390        |
| To Initiate Transcription, RNA Polymerase II Requires a Set of General Transcription Factors                            | 332        | Ribozymes Can Be Produced in the Laboratory                                                                                    | 390        |

|                                                                                                                |            |                                                                                                            |            |
|----------------------------------------------------------------------------------------------------------------|------------|------------------------------------------------------------------------------------------------------------|------------|
| RNA Can Both Store Information and Catalyze Chemical Reactions                                                 | 391        | Transcription Circuits Allow the Cell to Carry Out Logic Operations                                        | 433        |
| How Did Protein Synthesis Evolve?                                                                              | 392        | Summary                                                                                                    | 434        |
| All Present-Day Cells Use DNA as Their Hereditary Material                                                     | 393        | <b>MECHANISMS THAT REINFORCE CELL MEMORY</b>                                                               |            |
| Summary                                                                                                        | 393        | IN PLANTS AND ANIMALS                                                                                      | 435        |
| Problems                                                                                                       | 394        | Patterns of DNA Methylation Can Be Inherited When Vertebrate Cells Divide                                  | 435        |
| References                                                                                                     | 395        | CG-Rich Islands Are Associated with Many Genes in Mammals                                                  | 436        |
|                                                                                                                |            | Genomic Imprinting Is Based on DNA Methylation                                                             | 438        |
|                                                                                                                |            | A Chromosome-wide Alteration in Chromatin Structure Can Be Inherited                                       | 440        |
| <b>Chapter 7 Control of Gene Expression</b>                                                                    | <b>397</b> | The Mammalian X-Inactivation in Females Is Triggered by the Synthesis of a Long Noncoding RNA              | 442        |
| <b>AN OVERVIEW OF GENE CONTROL</b>                                                                             | 397        | Stable Patterns of Gene Expression Can Be Transmitted to Daughter Cells                                    | 443        |
| The Different Cell Types of a Multicellular Organism Contain the Same DNA                                      | 397        | Summary                                                                                                    | 445        |
| Different Cell Types Synthesize Different Sets of RNAs and Proteins                                            | 398        | <b>POST-TRANSCRIPTIONAL CONTROLS</b>                                                                       | 445        |
| The Spectrum of mRNAs Present in a Cell Can Be Used to Accurately Identify the Cell Type                       | 400        | Transcription Attenuation Causes the Premature Termination of Some RNA Molecules                           | 445        |
| External Signals Can Cause a Cell to Change the Expression of Its Genes                                        | 400        | Riboswitches Probably Represent Ancient Forms of Gene Control                                              | 446        |
| Gene Expression Can Be Regulated at Many of the Steps in the Pathway from DNA to RNA to Protein                | 401        | Alternative RNA Splicing Can Produce Different Forms of a Protein from the Same Gene                       | 446        |
| Summary                                                                                                        | 402        | The Definition of a Gene Has Been Modified Since the Discovery of Alternative RNA Splicing                 | 448        |
| <b>CONTROL OF TRANSCRIPTION BY SEQUENCE-SPECIFIC DNA-BINDING PROTEINS</b>                                      | 402        | Back Splicing Can Produce Circular RNA Molecules                                                           | 449        |
| The Sequence of Nucleotides in the DNA Double Helix Can Be Read by Proteins                                    | 402        | A Change in the Site of RNA Transcript Cleavage and Poly-A Addition Can Change the C-terminus of a Protein | 449        |
| Transcription Regulators Contain Structural Motifs That Can Read DNA Sequences                                 | 403        | Nucleotides in mRNA Can Be Covalently Modified                                                             | 450        |
| Dimerization of Transcription Regulators Increases Their Affinity and Specificity for DNA                      | 406        | RNA Editing Can Change the Meaning of the RNA Message                                                      | 451        |
| Many Transcription Regulators Bind Cooperatively to DNA                                                        | 407        | The Human AIDS Virus Illustrates How RNA Transport from the Nucleus Can Be Regulated                       | 452        |
| Nucleosome Structure Promotes Cooperative Binding of Transcription Regulators                                  | 408        | mRNAs Can Be Localized to Specific Regions of the Cytosol                                                  | 453        |
| DNA-Binding by Transcription Regulators Is Dynamic                                                             | 409        | Untranslated Regions of mRNAs Control Their Translation                                                    | 456        |
| Summary                                                                                                        | 410        | The Phosphorylation of an Initiation Factor Regulates Protein Synthesis Globally                           | 457        |
| <b>TRANSCRIPTION REGULATORS SWITCH GENES ON AND OFF</b>                                                        | 410        | Initiation at AUG Codons Upstream of the Translation Start Can Regulate Eukaryotic Translation Initiation  | 458        |
| The Tryptophan Repressor Switches Genes Off                                                                    | 410        | Internal Ribosome Entry Sites Also Provide Opportunities for Translational Control                         | 458        |
| Repressors Turn Genes Off and Activators Turn Them On                                                          | 411        | Changes in mRNA Stability Can Control Gene Expression                                                      | 459        |
| Both an Activator and a Repressor Control the <i>Lac</i> Operon                                                | 412        | Regulation of mRNA Stability Involves P-bodies and Stress Granules                                         | 461        |
| DNA Looping Can Occur During Bacterial Gene Regulation                                                         | 412        | Summary                                                                                                    | 462        |
| Complex Switches Control Gene Transcription in Eukaryotes                                                      | 414        | <b>REGULATION OF GENE EXPRESSION BY NONCODING RNAs</b>                                                     | 462        |
| A Eukaryotic Gene Control Region Includes Many <i>cis</i> -Regulatory Sequences                                | 414        | Small Noncoding RNA Transcripts Regulate Many Animal and Plant Genes Through RNA Interference              | 462        |
| Eukaryotic Transcription Regulators Work in Groups                                                             | 415        | miRNAs Regulate mRNA Translation and Stability                                                             | 463        |
| Activator Proteins Promote the Assembly of RNA Polymerase at the Start Point of Transcription                  | 416        | RNA Interference Also Serves as a Cell Defense Mechanism                                                   | 464        |
| Eukaryotic Transcription Activators Direct the Modification of Local Chromatin Structure                       | 417        | RNA Interference Can Direct Heterochromatin Formation                                                      | 465        |
| Some Transcription Activators Work by Releasing Paused RNA Polymerase                                          | 418        | piRNAs Protect the Germ Line from Transposable Elements                                                    | 466        |
| Transcription Activators Work Synergistically                                                                  | 419        | RNA Interference Has Become a Powerful Experimental Tool                                                   | 467        |
| Condensate Formation Likely Increases the Efficiency of Transcription Initiation                               | 420        | Cells Have Additional Mechanisms to Hold Transposons and Integrated Viral Genomes in Check                 | 467        |
| Eukaryotic Transcription Repressors Can Inhibit Transcription in Several Ways                                  | 420        | Bacteria Use Small Noncoding RNAs to Protect Themselves from Viruses                                       | 468        |
| Insulator DNA Sequences Prevent Eukaryotic Transcription Regulators from Influencing Distant Genes             | 422        | Long Noncoding RNAs Have Diverse Functions in the Cell                                                     | 469        |
| Summary                                                                                                        | 422        | Summary                                                                                                    | 471        |
|                                                                                                                |            | Problems                                                                                                   | 472        |
|                                                                                                                |            | References                                                                                                 | 474        |
| <b>MOLECULAR GENETIC MECHANISMS THAT CREATE AND MAINTAIN SPECIALIZED CELL TYPES</b>                            | <b>423</b> | <b>Chapter 8 Analyzing Cells, Molecules, and Systems</b>                                                   | <b>475</b> |
| Complex Genetic Switches That Regulate <i>Drosophila</i> Development Are Built Up from Smaller Modules         | 423        | <b>ISOLATING CELLS AND GROWING THEM IN CULTURE</b>                                                         | 476        |
| The <i>Drosophila Eve</i> Gene Is Regulated by Combinatorial Controls                                          | 424        | Cells Can Be Isolated from Tissues and Grown in Culture                                                    | 476        |
| Transcription Regulators Are Brought into Play by Extracellular Signals                                        | 426        | Eukaryotic Cell Lines Are a Widely Used Source of Homogeneous Cells                                        | 478        |
| Combinatorial Gene Control Creates Many Different Cell Types                                                   | 427        | Hybridoma Cell Lines Are Factories That Produce Monoclonal Antibodies                                      | 478        |
| Specialized Cell Types Can Be Experimentally Reprogrammed to Become Pluripotent Stem Cells                     | 428        | Summary                                                                                                    | 480        |
| Combinations of Master Transcription Regulators Specify Cell Types by Controlling the Expression of Many Genes | 429        | <b>PURIFYING PROTEINS</b>                                                                                  | 480        |
| Specialized Cells Must Rapidly Turn Some Genes On and Off                                                      | 430        | Cells Can Be Separated into Their Component Fractions                                                      | 480        |
| Differentiated Cells Maintain Their Identity                                                                   | 431        | Cell Extracts Provide Accessible Systems to Study Cell Functions                                           | 482        |
|                                                                                                                |            | Proteins Can Be Separated by Chromatography                                                                | 483        |

|                                                                                                                  |     |                                                                                                               |     |
|------------------------------------------------------------------------------------------------------------------|-----|---------------------------------------------------------------------------------------------------------------|-----|
| Immunoprecipitation Is a Rapid Affinity Purification Method                                                      | 486 | Expression of Individual Genes Can Be Measured Using                                                          |     |
| Genetically Engineered Tags Provide an Easy Way to Purify Proteins                                               | 486 | Quantitative RT-PCR                                                                                           | 536 |
| Purified Cell-free Systems Are Required for the Precise Dissection of Molecular Functions                        | 486 | Global Analysis of mRNAs by RNA-seq Provides a Snapshot of Gene Expression                                    | 536 |
| Summary                                                                                                          | 487 | Genome-wide Chromatin Immunoprecipitation Identifies Sites on the Genome Occupied by Transcription Regulators | 538 |
| <b>ANALYZING PROTEINS</b>                                                                                        | 487 | Ribosome Profiling Reveals Which mRNAs Are Being Translated in the Cell                                       | 538 |
| Proteins Can Be Separated by SDS Polyacrylamide-Gel Electrophoresis                                              | 487 | Recombinant DNA Methods Have Revolutionized Human Health                                                      | 539 |
| Two-dimensional Gel Electrophoresis Provides Greater Protein Separation                                          | 489 | Transgenic Plants Are Important for Agriculture                                                               | 540 |
| Specific Proteins Can Be Detected by Blotting with Antibodies                                                    | 490 | Summary                                                                                                       | 542 |
| Hydrodynamic Measurements Reveal the Size and Shape of a Protein Complex                                         | 490 | <b>MATHEMATICAL ANALYSIS OF CELL FUNCTION</b>                                                                 | 542 |
| Mass Spectrometry Provides a Highly Sensitive Method for Identifying Unknown Proteins                            | 491 | Regulatory Networks Depend on Molecular Interactions                                                          | 543 |
| Sets of Interacting Proteins Can Be Identified by Biochemical Methods                                            | 493 | Differential Equations Help Us Predict Transient Behavior                                                     | 545 |
| Optical Methods Can Monitor Protein Interactions                                                                 | 493 | Promoter Activity and Protein Degradation Affect the Rate of Change of Protein Concentration                  | 546 |
| Protein Structure Can Be Determined Using X-ray Diffraction                                                      | 494 | The Time Required to Reach Steady State Depends on Protein Lifetime                                           | 547 |
| NMR Can Be Used to Determine Protein Structure in Solution                                                       | 496 | Quantitative Methods Are Similar for Transcription Repressors and Activators                                  | 548 |
| Protein Sequence and Structure Provide Clues About Protein Function                                              | 497 | Negative Feedback Is a Powerful Strategy in Cell Regulation                                                   | 549 |
| Summary                                                                                                          | 498 | Delayed Negative Feedback Can Induce Oscillations                                                             | 549 |
| <b>ANALYZING AND MANIPULATING DNA</b>                                                                            | 498 | DNA Binding by a Repressor or an Activator Can Be Cooperative                                                 | 551 |
| Restriction Nucleases Cut Large DNA Molecules into Specific Fragments                                            | 498 | Positive Feedback Is Important for Switchlike Responses and Bistability                                       | 551 |
| Gel Electrophoresis Separates DNA Molecules of Different Sizes                                                   | 499 | Robustness Is an Important Characteristic of Biological Networks                                              | 553 |
| Purified DNA Molecules Can Be Specifically Labeled with Radioisotopes or Chemical Markers <i>in Vitro</i>        | 501 | Two Transcription Regulators That Bind to the Same Gene Promoter Can Exert Combinatorial Control              | 554 |
| Genes Can Be Cloned Using Bacteria                                                                               | 501 | An Incoherent Feed-forward Interaction Generates Pulses                                                       | 555 |
| An Entire Genome Can Be Represented in a DNA Library                                                             | 503 | A Coherent Feed-forward Interaction Detects Persistent Inputs                                                 | 556 |
| Hybridization Provides a Powerful but Simple Way to Detect Specific Nucleotide Sequences                         | 505 | The Same Network Can Behave Differently in Different Cells Because of Stochastic Effects                      | 557 |
| Genes Can Be Cloned <i>in Vitro</i> Using PCR                                                                    | 506 | Several Computational Approaches Can Be Used to Model the Reactions in Cells                                  | 557 |
| PCR Is Also Used for Diagnostic and Forensic Applications                                                        | 507 | Statistical Methods Are Critical for the Analysis of Biological Data                                          | 558 |
| PCR and Synthetic DNA Are Ideal Sources of Specific Gene Sequences for Cloning                                   | 510 | Summary                                                                                                       | 558 |
| DNA Cloning Allows Any Protein to Be Produced in Large Amounts                                                   | 511 | Problems                                                                                                      | 559 |
| DNA Can Be Sequenced Rapidly by Dideoxy Sequencing                                                               | 512 | References                                                                                                    | 561 |
| Next-Generation Sequencing Methods Have Revolutionized DNA and RNA Analysis                                      | 514 | <b>Chapter 9 Visualizing Cells and Their Molecules</b>                                                        | 563 |
| To Be Useful, Genome Sequences Must Be Annotated                                                                 | 516 | <b>LOOKING AT CELLS AND MOLECULES IN THE LIGHT MICROSCOPE</b>                                                 | 563 |
| Summary                                                                                                          | 518 | The Conventional Light Microscope Can Resolve Details 0.2 $\mu\text{m}$ Apart                                 | 564 |
| <b>STUDYING GENE FUNCTION AND EXPRESSION</b>                                                                     | 518 | Photon Noise Creates Additional Limits to Resolution When Light Levels Are Low                                | 567 |
| Classical Genetic Screens Identify Random Mutants with Specific Abnormalities                                    | 519 | Living Cells Are Seen Clearly in a Phase-Contrast or a Differential-Interference-Contrast Microscope          | 567 |
| Mutations Can Cause Loss or Gain of Protein Function                                                             | 522 | Images Can Be Enhanced and Analyzed by Digital Techniques                                                     | 568 |
| Complementation Tests Reveal Whether Two Mutations Are in the Same Gene or Different Genes                       | 523 | Intact Tissues Are Usually Fixed and Sectioned Before Microscopy                                              | 569 |
| Gene Products Can Be Ordered in Pathways by Epistasis Analysis                                                   | 523 | Specific Molecules Can Be Located in Cells by Fluorescence Microscopy                                         | 570 |
| Mutations Responsible for a Phenotype Can Be Identified Through DNA Analysis                                     | 524 | Antibodies Can Be Used to Detect Specific Proteins                                                            | 572 |
| Rapid and Cheap DNA Sequencing Has Revolutionized Human Genetic Studies                                          | 524 | Individual Proteins Can Be Fluorescently Tagged in Living Cells and Organisms                                 | 573 |
| Linked Blocks of Polymorphisms Have Been Passed Down from Our Ancestors                                          | 525 | Protein Dynamics Can Be Followed in Living Cells                                                              | 575 |
| Sequence Variants Can Aid the Search for Mutations Associated with Disease                                       | 526 | Fluorescent Biosensors Can Monitor Cell Signaling                                                             | 576 |
| Genomics Is Accelerating the Discovery of Rare Mutations That Predispose Us to Serious Disease                   | 527 | Imaging of Complex Three-dimensional Objects Is Possible with the Optical Microscope                          | 577 |
| The Cellular Functions of a Known Gene Can Be Studied with Genome Engineering                                    | 527 | The Confocal Microscope Produces Optical Sections by Excluding Out-of-Focus Light                             | 578 |
| Animals and Plants Can Be Genetically Altered                                                                    | 528 | Superresolution Fluorescence Techniques Can Overcome Diffraction-limited Resolution                           | 580 |
| The Bacterial CRISPR System Has Been Adapted to Edit Genomes in a Wide Variety of Species                        | 530 | Single-Molecule Localization Microscopy Also Delivers Superresolution                                         | 583 |
| Large Collections of Engineered Mutations Provide a Tool for Examining the Function of Every Gene in an Organism | 531 | Expanding the Specimen Can Offer Higher Resolution, but with a Conventional Microscope                        | 585 |
| RNA Interference Is a Simple and Rapid Way to Test Gene Function                                                 | 533 | Large Multicellular Structures Can Be Imaged Over Time                                                        | 586 |
| Reporter Genes Reveal When and Where a Gene Is Expressed                                                         | 534 | Single Molecules Can Be Visualized by Total Internal Reflection Fluorescence Microscopy                       | 587 |
| <i>In Situ</i> Hybridization Can Reveal the Location of mRNAs and Noncoding RNAs                                 | 535 | Summary                                                                                                       | 588 |

|                                                                                                                               |            |                                                                                                                                                  |            |
|-------------------------------------------------------------------------------------------------------------------------------|------------|--------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| <b>LOOKING AT CELLS AND MOLECULES IN THE ELECTRON MICROSCOPE</b>                                                              | <b>588</b> | A P-type ATPase Pumps $\text{Ca}^{2+}$ into the Sarcoplasmic Reticulum in Muscle Cells                                                           | <b>647</b> |
| The Electron Microscope Resolves the Fine Structure of the Cell                                                               | 588        | The Plasma Membrane $\text{Na}^{+}$ - $\text{K}^{+}$ Pump Establishes $\text{Na}^{+}$ and $\text{K}^{+}$ Gradients Across the Plasma Membrane    | 648        |
| Biological Specimens Require Special Preparation for Electron Microscopy                                                      | 589        | ABC Transporters Constitute the Largest Family of Membrane Transport Proteins                                                                    | 649        |
| Heavy Metals Can Provide Additional Contrast                                                                                  | 590        | Summary                                                                                                                                          | 651        |
| Images of Surfaces Can Be Obtained by Scanning Electron Microscopy                                                            | 591        | <b>CHANNELS AND THE ELECTRICAL PROPERTIES OF MEMBRANES</b>                                                                                       | <b>651</b> |
| Electron Microscope Tomography Allows the Molecular Architecture of Cells to Be Seen in Three Dimensions                      | 593        | Aquaporins Are Permeable to Water but Impermeable to Ions                                                                                        | 652        |
| Cryo-electron Microscopy Can Determine Molecular Structures at Atomic Resolution                                              | 595        | Ion Channels Are Ion-selective and Fluctuate Between Open and Closed States                                                                      | 653        |
| Light Microscopy and Electron Microscopy Are Mutually Beneficial                                                              | 597        | The Membrane Potential in Animal Cells Depends Mainly on $\text{K}^{+}$ Leak Channels and the $\text{K}^{+}$ Gradient Across the Plasma Membrane | 655        |
| Using Microscopy to Study Cells Always Involves Trade-Offs                                                                    | 598        | The Resting Potential Decays Only Slowly When the $\text{Na}^{+}$ - $\text{K}^{+}$ Pump Is Stopped                                               | 655        |
| Summary                                                                                                                       | 599        | The Three-dimensional Structure of a Bacterial $\text{K}^{+}$ Channel Shows How an Ion Channel Can Work                                          | 657        |
| Problems                                                                                                                      | 600        | Mechanosensitive Channels Allow Cells to Sense Their Physical Environment                                                                        | 659        |
| References                                                                                                                    | 601        | The Function of a Neuron Depends on Its Elongated Structure                                                                                      | 661        |
| <b>Chapter 10 Membrane Structure</b>                                                                                          | <b>603</b> | Voltage-gated Cation Channels Generate Action Potentials in Electrically Excitable Cells                                                         | 662        |
| <b>THE LIPID BILAYER</b>                                                                                                      | <b>604</b> | Myelination Increases the Speed and Efficiency of Action Potential Propagation in Nerve Cells                                                    | 666        |
| Glycerophospholipids, Sphingolipids, and Sterols Are the Major Lipids in Cell Membranes                                       | 605        | Patch-Clamp Recording Indicates That Individual Ion Channels Open in an All-or-Nothing Fashion                                                   | 666        |
| Phospholipids Spontaneously Form Bilayers                                                                                     | 606        | Voltage-gated Cation Channels Are Evolutionarily and Structurally Related                                                                        | 668        |
| The Lipid Bilayer Is a Two-dimensional Fluid                                                                                  | 608        | Different Neuron Types Display Characteristic Stable Firing Properties                                                                           | 668        |
| The Fluidity of a Lipid Bilayer Depends on Its Composition                                                                    | 609        | Transmitter-gated Ion Channels Convert Chemical Signals into Electrical Ones at Chemical Synapses                                                | 669        |
| Despite Their Fluidity, Lipid Bilayers Can Form Domains of Different Compositions                                             | 610        | Chemical Synapses Can Be Excitatory or Inhibitory                                                                                                | 670        |
| Lipid Droplets Are Surrounded by a Phospholipid Monolayer                                                                     | 611        | The Acetylcholine Receptors at the Neuromuscular Junction Are Excitatory Transmitter-gated Cation Channels                                       | 671        |
| The Asymmetry of the Lipid Bilayer Is Functionally Important                                                                  | 612        | Neurons Contain Many Types of Transmitter-gated Channels                                                                                         | 672        |
| Glycolipids Are Found on the Surface of All Eukaryotic Plasma Membranes                                                       | 613        | Many Psychoactive Drugs Act at Synapses                                                                                                          | 673        |
| Summary                                                                                                                       | 614        | Neuromuscular Transmission Involves the Sequential Activation of Five Different Sets of Ion Channels                                             | 673        |
| <b>MEMBRANE PROTEINS</b>                                                                                                      | <b>615</b> | Single Neurons Are Complex Computation Devices                                                                                                   | 674        |
| Membrane Proteins Can Be Associated with the Lipid Bilayer in Various Ways                                                    | 615        | Neuronal Computation Requires a Combination of at Least Three Kinds of $\text{K}^{+}$ Channels                                                   | 675        |
| Lipid Anchors Control the Membrane Localization of Some Signaling Proteins                                                    | 616        | Long-term Potentiation in the Mammalian Hippocampus Depends on $\text{Ca}^{2+}$ Entry Through NMDA-Receptor Channels                             | 677        |
| In Most Transmembrane Proteins, the Polypeptide Chain Crosses the Lipid Bilayer in an $\alpha$ -Helical Conformation          | 617        | The Use of Channelrhodopsins Has Revolutionized the Study of Neural Circuits                                                                     | 678        |
| Transmembrane $\alpha$ Helices Often Interact with One Another                                                                | 619        | Summary                                                                                                                                          | 679        |
| Some $\beta$ Barrels Form Large Channels                                                                                      | 619        | Problems                                                                                                                                         | 680        |
| Many Membrane Proteins Are Glycosylated                                                                                       | 621        | References                                                                                                                                       | 681        |
| Membrane Proteins Can Be Solubilized and Purified in Detergents                                                               | 622        | <b>Chapter 12 Intracellular Organization and Protein Sorting</b>                                                                                 | <b>683</b> |
| Bacteriorhodopsin Is a Light-driven Proton ( $\text{H}^{+}$ ) Pump That Traverses the Lipid Bilayer as Seven $\alpha$ Helices | 625        | <b>THE COMPARTMENTALIZATION OF CELLS</b>                                                                                                         | <b>683</b> |
| Membrane Proteins Often Function as Large Complexes                                                                           | 627        | All Eukaryotic Cells Have the Same Basic Set of Membrane-enclosed Organelles                                                                     | 683        |
| Many Membrane Proteins Diffuse in the Plane of the Membrane                                                                   | 627        | Evolutionary Origins Explain the Topological Relationships of Organelles                                                                         | 686        |
| Cells Can Confine Proteins and Lipids to Specific Domains Within a Membrane                                                   | 629        | Macromolecules Can Be Segregated Without a Surrounding Membrane                                                                                  | 688        |
| The Cortical Cytoskeleton Gives Membranes Mechanical Strength and Restricts Membrane Protein Diffusion                        | 630        | Multivalent Interactions Mediate Formation of Biomolecular Condensates                                                                           | 690        |
| Membrane-bending Proteins Deform Bilayers                                                                                     | 632        | Biomolecular Condensates Create Biochemical Factories                                                                                            | 690        |
| Summary                                                                                                                       | 633        | Biomolecular Condensates Form and Disassemble in Response to Need                                                                                | 693        |
| Problems                                                                                                                      | 634        | Proteins Can Move Between Compartments in Different Ways                                                                                         | 694        |
| References                                                                                                                    | 635        | Sorting Signals and Sorting Receptors Direct Proteins to the Correct Cell Address                                                                | 695        |
| <b>Chapter 11 Small-Molecule Transport and Electrical Properties of Membranes</b>                                             | <b>637</b> | Construction of Most Organelles Requires Information in the Organelle Itself                                                                     | 697        |
| <b>PRINCIPLES OF MEMBRANE TRANSPORT</b>                                                                                       | <b>637</b> | Summary                                                                                                                                          | 697        |
| Protein-free Lipid Bilayers Are Impermeable to Ions                                                                           | 638        |                                                                                                                                                  |            |
| There Are Two Main Classes of Membrane Transport Proteins: Transporters and Channels                                          | 638        |                                                                                                                                                  |            |
| Active Transport Is Mediated by Transporters Coupled to an Energy Source                                                      | 639        |                                                                                                                                                  |            |
| Summary                                                                                                                       | 640        |                                                                                                                                                  |            |
| <b>TRANSPORTERS AND ACTIVE MEMBRANE TRANSPORT</b>                                                                             | <b>640</b> |                                                                                                                                                  |            |
| Active Transport Can Be Driven by Ion-Concentration Gradients                                                                 | 642        |                                                                                                                                                  |            |
| Transporters in the Plasma Membrane Regulate Cytosolic pH                                                                     | 644        |                                                                                                                                                  |            |
| An Asymmetric Distribution of Transporters in Epithelial Cells Underlies the Transcellular Transport of Solutes               | 645        |                                                                                                                                                  |            |
| There Are Three Classes of ATP-driven Pumps                                                                                   | 646        |                                                                                                                                                  |            |

|                                                                                                                         |            |                                                                                                                      |            |
|-------------------------------------------------------------------------------------------------------------------------|------------|----------------------------------------------------------------------------------------------------------------------|------------|
| <b>THE ENDOPLASMIC RETICULUM</b>                                                                                        | <b>698</b> | The Assembly of a Clathrin Coat Drives Vesicle Formation                                                             | 752        |
| The ER Is Structurally and Functionally Diverse                                                                         | 698        | Adaptor Proteins Select Cargo into Clathrin-coated Vesicles                                                          | 753        |
| Signal Sequences Were First Discovered in Proteins Imported into the Rough ER                                           | 701        | Phosphoinositides Mark Organelles and Membrane Domains                                                               | 754        |
| A Signal-Recognition Particle (SRP) Directs the ER Signal Sequence to a Specific Receptor at the ER                     | 702        | Membrane-bending Proteins Help Deform the Membrane During Vesicle Formation                                          | 755        |
| The Polypeptide Chain Passes Through a Signal Sequence-gated Aqueous Channel in the Translocator                        | 705        | Cytoplasmic Proteins Regulate the Pinching off and Uncoating of Coated Vesicles                                      | 756        |
| Translocation Across the ER Membrane Does Not Always Require Ongoing Polypeptide Chain Elongation                       | 707        | Monomeric GTPases Control Coat Assembly                                                                              | 756        |
| Transmembrane Proteins Contain Hydrophobic Segments That Are Recognized Like Signal Sequences                           | 709        | Coat-recruitment GTPases Participate in Coat Disassembly                                                             | 758        |
| Hydrophobic Segments of Multipass Transmembrane Proteins Are Interpreted Contextually to Determine Their Orientation    | 710        | The Shape and Size of Transport Vesicles Are Diverse                                                                 | 759        |
| Some Proteins Are Integrated into the ER Membrane by a Post-translational Mechanism                                     | 711        | Rab Proteins Guide Transport Vesicles to Their Target Membrane                                                       | 760        |
| Some Membrane Proteins Acquire a Covalently Attached Glycosylphosphatidylinositol (GPI) Anchor                          | 712        | Rab Proteins Create and Change the Identity of an Organelle                                                          | 761        |
| Translocated Polypeptide Chains Fold and Assemble in the Lumen of the Rough ER                                          | 712        | SNAREs Mediate Membrane Fusion                                                                                       | 762        |
| Most Proteins Synthesized in the Rough ER Are Glycosylated by the Addition of a Common <i>N</i> -Linked Oligosaccharide | 714        | Interacting SNAREs Need to Be Pried Apart Before They Can Function Again                                             | 763        |
| Oligosaccharides Are Used as Tags to Mark the State of Protein Folding                                                  | 715        | Viruses Encode Specialized Membrane Fusion Proteins Needed for Cell Entry                                            | 764        |
| Improperly Folded Proteins Are Exported from the ER and Degraded in the Cytosol                                         | 716        | Summary                                                                                                              | 764        |
| Misfolded Proteins in the ER Activate an Unfolded Protein Response                                                      | 717        | <b>TRANSPORT FROM THE ENDOPLASMIC RETICULUM THROUGH THE GOLGI APPARATUS</b>                                          | <b>765</b> |
| The ER Assembles Most Lipid Bilayers                                                                                    | 720        | Proteins Leave the ER in COPII-coated Transport Vesicles                                                             | 765        |
| Membrane Contact Sites Between the ER and Other Organelles Facilitate Selective Lipid Transfer                          | 722        | Only Proteins That Are Properly Folded and Assembled Can Leave the ER                                                | 766        |
| Summary                                                                                                                 | 723        | Vesicular Tubular Clusters Mediate Transport from the ER to the Golgi Apparatus                                      | 766        |
| <b>PEROXISOMES</b>                                                                                                      | <b>723</b> | The Retrieval Pathway to the ER Uses Sorting Signals                                                                 | 768        |
| Peroxisomes Use Molecular Oxygen and Hydrogen Peroxide to Perform Oxidation Reactions                                   | 724        | Many Proteins Are Selectively Retained in the Compartments in Which They Function                                    | 768        |
| Short Signal Sequences Direct the Import of Proteins into Peroxisomes                                                   | 724        | The Golgi Apparatus Consists of an Ordered Series of Compartments                                                    | 769        |
| Summary                                                                                                                 | 726        | Oligosaccharide Chains Are Processed in the Golgi Apparatus                                                          | 771        |
| <b>THE TRANSPORT OF PROTEINS INTO MITOCHONDRIA AND CHLOROPLASTS</b>                                                     | <b>726</b> | Proteoglycans Are Assembled in the Golgi Apparatus                                                                   | 772        |
| Translocation into Mitochondria Depends on Signal Sequences and Protein Translocators                                   | 727        | What Is the Purpose of Glycosylation?                                                                                | 773        |
| Mitochondrial Proteins Are Imported Post-translationally as Unfolded Polypeptide Chains                                 | 728        | Transport Through the Golgi Apparatus Occurs by Multiple Mechanisms                                                  | 774        |
| Protein Import Is Powered by ATP Hydrolysis, a Membrane Potential, and Redox Potential                                  | 730        | Golgi Matrix Proteins Help Organize the Stack                                                                        | 775        |
| Transport into the Inner Mitochondrial Membrane Occurs Via Several Routes                                               | 731        | Summary                                                                                                              | 776        |
| Bacteria and Mitochondria Use Similar Mechanisms to Insert $\beta$ Barrels into Their Outer Membrane                    | 733        | <b>TRANSPORT FROM THE TRANS GOLGI NETWORK TO THE CELL EXTERIOR AND ENDOSOMES</b>                                     | <b>776</b> |
| Two Signal Sequences Direct Proteins to the Thylakoid Membrane in Chloroplasts                                          | 733        | Many Proteins and Lipids Are Carried Automatically from the <i>Trans</i> Golgi Network to the Cell Surface           | 777        |
| Summary                                                                                                                 | 735        | A Mannose 6-Phosphate Receptor Sorts Lysosomal Hydrolases in the <i>Trans</i> Golgi Network                          | 777        |
| <b>THE TRANSPORT OF MOLECULES BETWEEN THE NUCLEUS AND THE CYTOSOL</b>                                                   | <b>735</b> | Defects in the GlcNAc Phosphotransferase Cause a Lysosomal Storage Disease in Humans                                 | 779        |
| Nuclear Pore Complexes Perforate the Nuclear Envelope                                                                   | 736        | Secretory Vesicles Bud from the <i>Trans</i> Golgi Network                                                           | 780        |
| Nuclear Localization Signals Direct Proteins to the Nucleus                                                             | 738        | Precursors of Secretory Proteins Are Proteolytically Processed During the Formation of Secretory Vesicles            | 781        |
| Nuclear Import Receptors Bind to Both Nuclear Localization Signals and NPC Proteins                                     | 739        | Secretory Vesicles Wait Near the Plasma Membrane Until Signaled to Release Their Contents                            | 782        |
| The Ran GTPase Imposes Directionality on Nuclear Import Through NPCs                                                    | 740        | For Rapid Exocytosis, Synaptic Vesicles Are Primed at the Presynaptic Plasma Membrane                                | 782        |
| Nuclear Export Works Like Nuclear Import, but in Reverse                                                                | 741        | Synaptic Vesicles Can Be Recycled Locally After Exocytosis                                                           | 783        |
| Transport Through NPCs Can Be Regulated by Controlling Access to the Transport Machinery                                | 742        | Secretory Vesicle Membrane Components Are Quickly Removed from the Plasma Membrane                                   | 784        |
| The Nuclear Envelope Disassembles and Reassembles During Mitosis                                                        | 743        | Some Regulated Exocytosis Events Serve to Enlarge the Plasma Membrane                                                | 785        |
| Summary                                                                                                                 | 745        | Polarized Cells Direct Proteins from the <i>Trans</i> Golgi Network to the Appropriate Domain of the Plasma Membrane | 786        |
| Problems                                                                                                                | 746        | Summary                                                                                                              | 787        |
| References                                                                                                              | 748        | <b>TRANSPORT INTO THE CELL FROM THE PLASMA MEMBRANE: ENDOCYTOSIS</b>                                                 | <b>788</b> |
| <b>Chapter 13 Intracellular Membrane Traffic</b>                                                                        | <b>749</b> | Pinocytic Vesicles Form from Coated Pits in the Plasma Membrane                                                      | 789        |
| <b>MECHANISMS OF MEMBRANE TRANSPORT AND COMPARTMENT IDENTITY</b>                                                        | <b>751</b> | Not All Membrane Invaginations and Pinocytic Vesicles Are Clathrin Coated                                            | 789        |
| There Are Various Types of Coated Vesicles                                                                              | 751        | Cells Use Receptor-mediated Endocytosis to Import Selected Extracellular Macromolecules                              | 791        |
|                                                                                                                         |            | Specific Proteins Are Retrieved from Early Endosomes and Returned to the Plasma Membrane                             | 792        |
|                                                                                                                         |            | Recycling Endosomes Regulate Plasma Membrane Composition                                                             | 793        |
|                                                                                                                         |            | Plasma Membrane Signaling Receptors Are Down-regulated by Degradation in Lysosomes                                   | 794        |

|                                                                                                                        |     |                                                                                                                                                                  |     |
|------------------------------------------------------------------------------------------------------------------------|-----|------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Early Endosomes Mature into Late Endosomes                                                                             | 795 | CHLOROPLASTS AND PHOTOSYNTHESIS                                                                                                                                  | 843 |
| ESCRT Protein Complexes Mediate the Formation of Intraluminal Vesicles in Multivesicular Bodies                        | 796 | Chloroplasts Resemble Mitochondria but Have a Separate Thylakoid Compartment                                                                                     | 843 |
| Summary                                                                                                                | 798 | Chloroplasts Capture Energy from Sunlight and Use It to Fix Carbon                                                                                               | 844 |
| THE DEGRADATION AND RECYCLING OF MACROMOLECULES IN LYSOSOMES                                                           | 798 | Carbon Fixation Uses ATP and NADPH to Convert CO <sub>2</sub> into Sugars                                                                                        | 845 |
| Lysosomes Are the Principal Sites of Intracellular Digestion                                                           | 798 | Carbon Fixation in Some Plants Is Compartmentalized to Facilitate Growth at Low CO <sub>2</sub> Concentrations                                                   | 846 |
| Lysosomes Are Heterogeneous                                                                                            | 799 | The Sugars Generated by Carbon Fixation Can Be Stored as Starch or Consumed to Produce ATP                                                                       | 849 |
| Plant and Fungal Vacuoles Are Remarkably Versatile Lysosomes                                                           | 800 | The Thylakoid Membranes of Chloroplasts Contain the Protein Complexes Required for Photosynthesis and ATP Generation                                             | 849 |
| Multiple Pathways Deliver Materials to Lysosomes                                                                       | 801 | Chlorophyll-Protein Complexes Can Transfer Either Excitation Energy or Electrons                                                                                 | 850 |
| Cells Can Acquire Nutrients from the Extracellular Fluid by Macropinocytosis                                           | 802 | A Photosystem Contains Chlorophylls in Antennae and a Reaction Center                                                                                            | 851 |
| Specialized Phagocytic Cells Can Ingest Large Particles                                                                | 802 | The Thylakoid Membrane Contains Two Different Photosystems Working in Series                                                                                     | 852 |
| Cargo Recognition by Cell-surface Receptors Initiates Phagocytosis                                                     | 803 | Photosystem II Uses a Manganese Cluster to Withdraw Electrons from Water                                                                                         | 853 |
| Autophagy Degrades Unwanted Proteins and Organelles                                                                    | 804 | The Cytochrome <i>b<sub>6</sub>-f</i> Complex Connects Photosystem II to Photosystem I                                                                           | 854 |
| The Rate of Nonselective Autophagy Is Regulated by Nutrient Availability                                               | 805 | Photosystem I Carries Out the Second Charge-Separation Step in the Z Scheme                                                                                      | 855 |
| A Family of Cargo-specific Receptors Mediates Selective Autophagy                                                      | 806 | The Chloroplast ATP Synthase Uses the Proton Gradient Generated by the Photosynthetic Light Reactions to Produce ATP                                             | 855 |
| Some Lysosomes and Multivesicular Bodies Undergo Exocytosis                                                            | 807 | The Proton-Motive Force for ATP Production in Mitochondria and Chloroplasts Is Essentially the Same                                                              | 856 |
| Summary                                                                                                                | 807 | Chemiosmotic Mechanisms Evolved in Stages By Providing an Inexhaustible Source of Reducing Power, Photosynthetic Bacteria Overcame a Major Evolutionary Obstacle | 857 |
| Problems                                                                                                               | 808 | The Photosynthetic Electron-Transport Chains of Cyanobacteria Produced Atmospheric Oxygen and Permitted New Life-Forms                                           | 857 |
| References                                                                                                             | 810 | Summary                                                                                                                                                          | 860 |
| Chapter 14 Energy Conversion and Metabolic Compartmentation: Mitochondria and Chloroplasts                             | 811 | THE GENETIC SYSTEMS OF MITOCHONDRIA AND CHLOROPLASTS                                                                                                             | 861 |
| THE MITOCHONDRION                                                                                                      | 813 | The Genetic Systems of Mitochondria and Chloroplasts Resemble Those of Prokaryotes                                                                               | 861 |
| The Mitochondrion Has an Outer Membrane and an Inner Membrane                                                          | 814 | Over Time, Mitochondria and Chloroplasts Have Exported Most of Their Genes to the Nucleus by Gene Transfer                                                       | 862 |
| Fission, Fusion, Distribution, and Degradation of Mitochondria                                                         | 815 | Mitochondria Have a Relaxed Codon Usage and Can Have a Variant Genetic Code                                                                                      | 864 |
| The Inner Membrane Cristae Contain the Machinery for Electron Transport and ATP Synthesis                              | 817 | Chloroplasts and Bacteria Share Many Striking Similarities                                                                                                       | 865 |
| The Citric Acid Cycle in the Matrix Produces NADH                                                                      | 817 | Organelle Genes Are Maternally Inherited in Animals and Plants                                                                                                   | 866 |
| Mitochondria Have Many Essential Roles in Cellular Metabolism                                                          | 818 | Mutations in Mitochondrial DNA Can Cause Severe Inherited Diseases                                                                                               | 866 |
| A Chemiosmotic Process Couples Oxidation Energy to ATP Production                                                      | 821 | Why Do Mitochondria and Chloroplasts Maintain a Costly Separate System for DNA Transcription and Translation?                                                    | 867 |
| The Energy Derived from Oxidation Is Stored as an Electrochemical Gradient                                             | 822 | Summary                                                                                                                                                          | 868 |
| Summary                                                                                                                | 823 | Problems                                                                                                                                                         | 869 |
| THE PROTON PUMPS OF THE ELECTRON-TRANSPORT CHAIN                                                                       | 823 | References                                                                                                                                                       | 871 |
| The Redox Potential Is a Measure of Electron Affinities                                                                | 823 | Chapter 15 Cell Signaling                                                                                                                                        | 873 |
| Electron Transfers Release Large Amounts of Energy                                                                     | 824 | PRINCIPLES OF CELL SIGNALING                                                                                                                                     | 873 |
| Transition Metal Ions and Quinones Accept and Release Electrons Readily                                                | 824 | Extracellular Signals Can Act Over Short or Long Distances                                                                                                       | 874 |
| NADH Transfers Its Electrons to Oxygen Through Three Large Enzyme Complexes Embedded in the Inner Membrane             | 827 | Extracellular Signal Molecules Bind to Specific Receptors                                                                                                        | 875 |
| The NADH Dehydrogenase Complex Contains Separate Modules for Electron Transport and Proton Pumping                     | 828 | Each Cell Is Programmed to Respond to Specific Combinations of Extracellular Signals                                                                             | 876 |
| Cytochrome c Reductase Takes Up and Releases Protons on Opposite Sides of the Crista Membrane, Thereby Pumping Protons | 829 | There Are Three Major Classes of Cell-Surface Receptor Proteins                                                                                                  | 878 |
| The Cytochrome c Oxidase Complex <i>Pumps</i> Protons and Reduces O <sub>2</sub> Using a Catalytic Iron-Copper Center  | 831 | Cell-Surface Receptors Relay Signals Via Intracellular Signaling Molecules                                                                                       | 879 |
| Succinate Dehydrogenase Acts in Both the Electron-Transport Chain and the Citric Acid Cycle                            | 832 | Intracellular Signals Must Be Specific and Robust in a Noisy Cytoplasm                                                                                           | 881 |
| The Respiratory Chain Forms a Supercomplex in the Crista Membrane                                                      | 833 | Intracellular Signaling Complexes Form at Activated Cell-Surface Receptors                                                                                       | 882 |
| Protons Can Move Rapidly Through Proteins Along Predefined Pathways                                                    | 834 | Modular Interaction Domains Mediate Interactions Between Intracellular Signaling Proteins                                                                        | 883 |
| Summary                                                                                                                | 835 | The Relationship Between Signal and Response Varies in Different Signaling Pathways                                                                              | 885 |
| ATP PRODUCTION IN MITOCHONDRIA                                                                                         | 835 | The Speed of a Response Depends on the Turnover of Signaling Molecules                                                                                           | 886 |
| The Large Negative Value of $\Delta G$ for ATP Hydrolysis Makes ATP Useful to the Cell                                 | 835 |                                                                                                                                                                  |     |
| The ATP Synthase Is a Nanomachine That Produces ATP by Rotary Catalysis                                                | 837 |                                                                                                                                                                  |     |
| Proton-driven Turbines Are Ancient and Critical for Energy Conversion                                                  | 839 |                                                                                                                                                                  |     |
| Mitochondrial Cristae Help to Make ATP Synthesis Efficient                                                             | 840 |                                                                                                                                                                  |     |
| Special Transport Proteins Move Solutes Through the Inner Membrane                                                     | 841 |                                                                                                                                                                  |     |
| Chemiosmotic Mechanisms First Arose in Bacteria                                                                        | 842 |                                                                                                                                                                  |     |
| Summary                                                                                                                | 842 |                                                                                                                                                                  |     |

|                                                                                                                      |            |                                                                                                                |             |
|----------------------------------------------------------------------------------------------------------------------|------------|----------------------------------------------------------------------------------------------------------------|-------------|
| Cells Can Respond Abruptly to a Gradually Increasing Signal                                                          | 887        | <b>Chapter 16 The Cytoskeleton</b>                                                                             | <b>949</b>  |
| Positive Feedback Can Generate an All-or-None Response                                                               | 888        | <b>FUNCTION AND DYNAMICS OF THE CYTOSKELETON</b>                                                               | <b>949</b>  |
| Negative Feedback Is a Common Feature of Intracellular Signaling Systems                                             | 890        | Cytoskeletal Filaments Are Dynamic, but Can Nevertheless Form Stable Structures                                | 951         |
| Cells Can Adjust Their Sensitivity to a Signal                                                                       | 890        | The Cytoskeleton Determines Cellular Organization and Polarity                                                 | 952         |
| Summary                                                                                                              | 892        | Filaments Assemble from Protein Subunits That Impart Specific Physical and Dynamic Properties                  | 953         |
| <b>SIGNALING THROUGH G-PROTEIN-COUPLED RECEPTORS</b>                                                                 | <b>892</b> | Accessory Proteins and Motors Act on Cytoskeletal Filaments                                                    | 955         |
| Heterotrimeric G Proteins Relay Signals from GPCRs                                                                   | 893        | Molecular Motors Operate in a Cellular Environment Dominated by Brownian Motion                                | 956         |
| Some G Proteins Regulate the Production of Cyclic AMP                                                                | 895        | Summary                                                                                                        | 957         |
| Cyclic-AMP-dependent Protein Kinase (PKA) Mediates Most of the Effects of Cyclic AMP                                 | 896        | <b>ACTIN</b>                                                                                                   | <b>957</b>  |
| Some G Proteins Signal Via Phospholipids                                                                             | 898        | Actin Subunits Assemble Head-to-Tail to Create Flexible, Polar Filaments                                       | 958         |
| Ca <sup>2+</sup> Functions as a Ubiquitous Intracellular Mediator                                                    | 899        | Nucleation Is the Rate-limiting Step in the Formation of Actin Filaments                                       | 958         |
| Feedback Generates Ca <sup>2+</sup> Waves and Oscillations                                                           | 900        | Actin Filaments Have Two Distinct Ends That Grow at Different Rates                                            | 962         |
| Ca <sup>2+</sup> /Calmodulin-dependent Protein Kinases Mediate Many Responses to Ca <sup>2+</sup> Signals            | 902        | ATP Hydrolysis Within Actin Filaments Leads to Treadmilling at Steady State                                    | 962         |
| Some G Proteins Directly Regulate Ion Channels                                                                       | 904        | The Functions of Actin Filaments Are Inhibited by Both Polymer-stabilizing and Polymer-destabilizing Chemicals | 963         |
| Smell and Vision Depend on GPCRs That Regulate Ion Channels                                                          | 905        | Actin-binding Proteins Influence Filament Dynamics and Organization                                            | 964         |
| Nitric Oxide Gas Can Mediate Signaling Between Cells                                                                 | 908        | Actin Nucleation Is Tightly Regulated and Generates Branched or Straight Filaments                             | 964         |
| Second Messengers and Enzymatic Cascades Amplify Signals                                                             | 909        | Actin Filament Elongation Is Regulated by Monomer-binding Proteins                                             | 967         |
| GPCR Desensitization Depends on Receptor Phosphorylation                                                             | 909        | Actin Filament-binding Proteins Alter Filament Dynamics and Organization                                       | 968         |
| Summary                                                                                                              | 910        | Severing Proteins Regulate Actin Filament Depolymerization                                                     | 970         |
| <b>SIGNALING THROUGH ENZYME-COUPLED RECEPTORS</b>                                                                    | <b>911</b> | Bacteria Can Hijack the Host Actin Cytoskeleton                                                                | 971         |
| Activated Receptor Tyrosine Kinases (RTKs) Phosphorylate Themselves                                                  | 911        | Actin at the Cell Cortex Determines Cell Shape                                                                 | 971         |
| Phosphorylated Tyrosines on RTKs Serve as Docking Sites for Intracellular Signaling Proteins                         | 913        | Distinct Modes of Cell Migration Rely on the Actin Cytoskeleton                                                | 972         |
| Proteins with SH2 Domains Bind to Phosphorylated Tyrosines                                                           | 913        | Cells Migrating in Three Dimensions Can Navigate Around Barriers                                               | 974         |
| The Monomeric GTPase Ras Mediates Signaling by Most RTKs                                                             | 915        | Summary                                                                                                        | 975         |
| Ras Activates a MAP Kinase Signaling Module                                                                          | 916        | <b>MYOSIN AND ACTIN</b>                                                                                        | <b>976</b>  |
| Scaffold Proteins Reduce Cross-Talk Between Different MAP Kinase Modules                                             | 918        | Actin-based Motor Proteins Are Members of the Myosin Superfamily                                               | 976         |
| Rho Family GTPases Functionally Couple Cell-Surface Receptors to the Cytoskeleton                                    | 919        | Myosin Generates Force by Coupling ATP Hydrolysis to Conformational Changes                                    | 977         |
| PI 3-Kinase Produces Lipid Docking Sites in the Plasma Membrane                                                      | 920        | Sliding of Myosin II Along Actin Filaments Causes Muscles to Contract                                          | 977         |
| The PI-3-Kinase-Akt Signaling Pathway Stimulates Animal Cells to Survive and Grow                                    | 921        | A Sudden Rise in Cytosolic Ca <sup>2+</sup> Concentration Initiates Muscle Contraction                         | 981         |
| RTKs and GPCRs Activate Overlapping Signaling Pathways                                                               | 923        | Heart Muscle Is a Precisely Engineered Machine                                                                 | 984         |
| Some Enzyme-coupled Receptors Associate with Cytoplasmic Tyrosine Kinases                                            | 923        | Actin and Myosin Perform a Variety of Functions in Non-Muscle Cells                                            | 984         |
| Cytokine Receptors Activate the JAK-STAT Signaling Pathway                                                           | 924        | Summary                                                                                                        | 986         |
| Extracellular Signal Proteins of the TGF $\beta$ Superfamily Act Through Receptor Serine/Threonine Kinases and Smads | 926        | <b>MICROTUBULES</b>                                                                                            | <b>987</b>  |
| Summary                                                                                                              | 927        | Microtubules Are Hollow Tubes Made of Protofilaments                                                           | 988         |
| <b>ALTERNATIVE SIGNALING ROUTES IN GENE REGULATION</b>                                                               | <b>928</b> | Microtubules Undergo a Process Called Dynamic Instability                                                      | 988         |
| The Receptor Notch Is a Latent Transcription Regulator                                                               | 928        | Microtubule Functions Are Inhibited by Both Polymer-stabilizing and Polymer-destabilizing Drugs                | 991         |
| Wnt Proteins Activate Frizzled and Thereby Inhibit $\beta$ -Catenin Degradation                                      | 930        | A Protein Complex Containing $\gamma$ -Tubulin Nucleates Microtubules                                          | 991         |
| Hedgehog Proteins Initiate a Complex Signaling Pathway in the Primary Cilium                                         | 932        | The Centrosome Is a Prominent Microtubule Nucleation Site                                                      | 991         |
| Many Inflammatory and Stress Signals Act Through an NF $\kappa$ B-dependent Signaling Pathway                        | 934        | Microtubule Organization Varies Widely Among Cell Types                                                        | 993         |
| Nuclear Receptors Are Ligand-modulated Transcription Regulators                                                      | 935        | Microtubule-binding Proteins Modulate Filament Dynamics and Organization                                       | 995         |
| Circadian Clocks Use Negative Feedback Loops to Control Gene Expression                                              | 937        | Microtubule Plus End-binding Proteins Modulate Microtubule Dynamics and Attachments                            | 996         |
| Three Purified Proteins Can Reconstitute a Cyanobacterial Circadian Clock in a Test Tube                             | 938        | Tubulin-sequestering and Microtubule-severing Proteins Modulate Microtubule Dynamics                           | 998         |
| Summary                                                                                                              | 940        | Two Types of Motor Proteins Move Along Microtubules                                                            | 999         |
| <b>SIGNALING IN PLANTS</b>                                                                                           | <b>940</b> | Microtubules and Motors Move Organelles and Vesicles                                                           | 1002        |
| Multicellularity and Cell Communication Evolved Independently in Plants and Animals                                  | 940        | Motile Cilia and Flagella Are Built from Microtubules and Dyneins                                              | 1004        |
| Receptor Serine/Threonine Kinases Are the Largest Class of Cell-Surface Receptors in Plants                          | 941        | Primary Cilia Perform Important Signaling Functions in Animal Cells                                            | 1005        |
| Ethylene Blocks the Degradation of Specific Transcription Regulatory Proteins in the Nucleus                         | 941        | Summary                                                                                                        | 1006        |
| Regulated Positioning of Auxin Transporters Patterns Plant Growth                                                    | 943        | <b>INTERMEDIATE FILAMENTS AND OTHER CYTOSKELETAL POLYMERS</b>                                                  | <b>1007</b> |
| Phytochromes Detect Red Light, and Cryptochromes Detect Blue Light                                                   | 944        | Intermediate Filament Structure Depends on the Lateral Bundling and Twisting of Coiled-Coils                   | 1007        |
| Summary                                                                                                              | 945        | Intermediate Filaments Impart Mechanical Stability to Animal Cells                                             | 1009        |
| Problems                                                                                                             | 946        | Linker Proteins Connect Cytoskeletal Filaments and Bridge the Nuclear Envelope                                 | 1011        |
| References                                                                                                           | 948        |                                                                                                                |             |

|                                                                                                |             |                                                                                                                                     |             |
|------------------------------------------------------------------------------------------------|-------------|-------------------------------------------------------------------------------------------------------------------------------------|-------------|
| Septins Form Filaments That Contribute to Subcellular Organization                             | 1012        | Actin and Myosin II in the Contractile Ring Guide the Process of Cytokinesis                                                        | 1065        |
| Bacterial Cell Shape and Division Depend on Homologs of Eukaryotic Cytoskeletal Proteins       | 1013        | Local Activation of RhoA Triggers Assembly and Contraction of the Contractile Ring                                                  | 1065        |
| Summary                                                                                        | 1016        | The Microtubules of the Mitotic Spindle Determine the Plane of Animal Cell Division                                                 | 1066        |
| <b>CELL POLARITY AND COORDINATION OF THE CYTOSKELETON</b>                                      | 1016        | The Phragmoplast Guides Cytokinesis in Higher Plants                                                                                | 1068        |
| Cell Polarity Is Governed by Small GTPases in Budding Yeast                                    | 1016        | Membrane-enclosed Organelles Must Be Distributed to Daughter Cells During Cytokinesis                                               | 1069        |
| PAR Proteins Generate Anterior–Posterior Polarity in Embryos                                   | 1018        | Some Cells Reposition Their Spindle to Divide Asymmetrically                                                                        | 1069        |
| Conserved Complexes Polarize Epithelial Cells and Control Their Growth                         | 1019        | Mitosis Can Occur Without Cytokinesis                                                                                               | 1070        |
| Cell Migration Requires Dynamic Cell Polarity                                                  | 1020        | Summary                                                                                                                             | 1070        |
| External Signals Can Dictate the Direction of Cell Migration                                   | 1022        | <b>MEIOSIS</b>                                                                                                                      | 1071        |
| Communication Among Cytoskeletal Elements Supports Whole-Cell Polarity and Locomotion          | 1023        | Meiosis Includes Two Rounds of Chromosome Segregation                                                                               | 1071        |
| Summary                                                                                        | 1023        | Duplicated Homologs Pair During Meiotic Prophase                                                                                    | 1073        |
| Problems                                                                                       | 1024        | Homolog Pairing Culminates in the Formation of a Synaptonemal Complex                                                               | 1073        |
| References                                                                                     | 1025        | Homolog Segregation Depends on Several Unique Features of Meiosis I                                                                 | 1075        |
| <b>Chapter 17 The Cell Cycle</b>                                                               | <b>1027</b> | Crossing-Over Is Highly Regulated                                                                                                   | 1076        |
| <b>OVERVIEW OF THE CELL CYCLE</b>                                                              | 1027        | Meiosis Frequently Goes Wrong                                                                                                       | 1077        |
| The Eukaryotic Cell Cycle Usually Consists of Four Phases                                      | 1028        | Summary                                                                                                                             | 1077        |
| Cell-Cycle Control Is Similar in All Eukaryotes                                                | 1030        | <b>CONTROL OF CELL DIVISION AND CELL GROWTH</b>                                                                                     | 1077        |
| Cell-Cycle Progression Can Be Studied in Various Ways                                          | 1030        | Mitogens Stimulate Cell Division                                                                                                    | 1078        |
| Summary                                                                                        | 1031        | Cells Can Enter a Specialized Nondividing State                                                                                     | 1078        |
| <b>THE CELL-CYCLE CONTROL SYSTEM</b>                                                           | 1031        | Mitogens Stimulate G <sub>1</sub> -Cdk and G <sub>1</sub> /S-Cdk Activities                                                         | 1079        |
| The Cell-Cycle Control System Triggers the Major Events of the Cell Cycle                      | 1032        | DNA Damage Blocks Cell Division                                                                                                     | 1080        |
| The Cell-Cycle Control System Depends on Cyclically Activated Cyclin-dependent Protein Kinases | 1033        | Many Human Cells Have a Built-In Limitation on the Number of Times They Can Divide                                                  | 1082        |
| Protein Phosphatases Reverse the Effects of Cdks                                               | 1035        | Cell Proliferation Is Accompanied by Cell Growth                                                                                    | 1083        |
| Hundreds of Cdk Substrates Are Phosphorylated in a Defined Order                               | 1035        | Proliferating Cells Usually Coordinate Their Growth and Division                                                                    | 1084        |
| Positive Feedback Generates the Switchlike Behavior of Cell-Cycle Transitions                  | 1036        | Summary                                                                                                                             | 1084        |
| The Anaphase-promoting Complex/Cyclosome (APC/C) Triggers the Metaphase-to-Anaphase Transition | 1038        | Problems                                                                                                                            | 1085        |
| The G <sub>1</sub> Phase Is a Stable State of Cdk Inactivity                                   | 1040        | References                                                                                                                          | 1087        |
| The Cell-Cycle Control System Functions as a Linked Series of Biochemical Switches             | 1041        | <b>Chapter 18 Cell Death</b>                                                                                                        | <b>1089</b> |
| Summary                                                                                        | 1042        | Apoptosis Eliminates Unwanted Cells                                                                                                 | 1090        |
| <b>S PHASE</b>                                                                                 | 1042        | Apoptosis Depends on an Intracellular Proteolytic Cascade Mediated by Caspases                                                      | 1091        |
| S-Cdk Initiates DNA Replication Once Per Cell Cycle                                            | 1043        | Activation of Cell-Surface Death Receptors Initiates the Extrinsic Pathway of Apoptosis                                             | 1093        |
| Chromosome Duplication Requires Duplication of Chromatin Structure                             | 1045        | The Intrinsic Pathway of Apoptosis Depends on Proteins Released from Mitochondria                                                   | 1094        |
| Cohesins Hold Sister Chromatids Together                                                       | 1045        | Bcl2 Proteins Are the Critical Controllers of the Intrinsic Pathway of Apoptosis                                                    | 1095        |
| Summary                                                                                        | 1046        | An Inhibitor of Apoptosis (an IAP) and Two Anti-IAP Proteins Help Control Caspase Activation in the Cytosol of Some Mammalian Cells | 1098        |
| <b>MITOSIS</b>                                                                                 | 1046        | Extracellular Survival Factors Inhibit Apoptosis in Various Ways                                                                    | 1098        |
| M-Cdk and Other Protein Kinases Drive Entry into Mitosis                                       | 1047        | Healthy Neighbors Phagocytose and Digest Apoptotic Cells                                                                            | 1100        |
| Condensin Helps Configure Duplicated Chromosomes for Separation                                | 1047        | Either Excessive or Insufficient Apoptosis Can Contribute to Disease                                                                | 1100        |
| The Mitotic Spindle Is a Dynamic Microtubule-based Machine                                     | 1050        | Summary                                                                                                                             | 1102        |
| Microtubules Are Nucleated in Multiple Regions of the Spindle                                  | 1051        | Problems                                                                                                                            | 1103        |
| Microtubule Instability Increases Greatly in Mitosis                                           | 1052        | References                                                                                                                          | 1104        |
| Microtubule-based Motor Proteins Govern Spindle Assembly and Function                          | 1052        | <b>Chapter 19 Cell Junctions and the Extracellular Matrix</b>                                                                       | <b>1105</b> |
| Bipolar Spindle Assembly in Most Animal Cells Begins with Centrosome Duplication               | 1053        | <b>CELL–CELL JUNCTIONS</b>                                                                                                          | 1108        |
| Spindle Assembly in Animal Cells Requires Nuclear-Envelope Breakdown                           | 1054        | Cadherins Form a Diverse Family of Adhesion Molecules                                                                               | 1108        |
| Mitotic Chromosomes Promote Bipolar Spindle Assembly                                           | 1055        | Cadherins Mediate Homophilic Adhesion                                                                                               | 1108        |
| Kinetochores Attach Sister Chromatids to the Spindle                                           | 1056        | Cadherin-dependent Cell–Cell Adhesion Guides the Organization of Developing Tissues                                                 | 1110        |
| Bi-orientation Is Achieved by Trial and Error                                                  | 1057        | Assembly of Strong Cell–Cell Adhesions Requires Changes in the Actin Cytoskeleton                                                   | 1112        |
| Multiple Forces Act on Chromosomes in the Spindle                                              | 1059        | Catenins Link Classical Cadherins to the Actin Cytoskeleton                                                                         | 1113        |
| The APC/C Triggers Sister-Chromatid Separation and the Completion of Mitosis                   | 1060        | Adherens Junctions Respond to Tension from Inside and Outside the Tissue                                                            | 1113        |
| Unattached Chromosomes Block Sister-Chromatid Separation: The Spindle Assembly Checkpoint      | 1062        | Tissue Remodeling Depends on the Coordination of Actin-mediated Contraction with Cell–Cell Adhesion                                 | 1114        |
| Chromosomes Segregate in Anaphase A and B                                                      | 1062        | Desmosomes Give Epithelia Mechanical Strength                                                                                       | 1116        |
| Segregated Chromosomes Are Packaged in Daughter Nuclei at Telophase                            | 1063        | Tight Junctions Form a Seal Between Cells and a Fence Between Plasma Membrane Domains                                               | 1116        |
| Summary                                                                                        | 1064        |                                                                                                                                     |             |
| <b>CYTOKINESIS</b>                                                                             | 1064        |                                                                                                                                     |             |

|                                                                                                                |      |                                                                                                                           |      |
|----------------------------------------------------------------------------------------------------------------|------|---------------------------------------------------------------------------------------------------------------------------|------|
| Tight Junctions Contain Strands of Transmembrane Adhesion Proteins                                             | 1119 | Cancer Cells Contain Somatic Mutations                                                                                    | 1166 |
| Scaffold Proteins Organize Junctional Protein Complexes                                                        | 1120 | A Single Mutation Is Not Enough to Change a Normal Cell into a Cancer Cell                                                | 1166 |
| Gap Junctions Couple Cells Both Electrically and Metabolically                                                 | 1121 | Many Cancers Develop Gradually Through Successive Rounds of Random Inherited Change Followed by Natural Selection         | 1167 |
| A Gap-Junction Connexon Is Made of Six Transmembrane Connexin Subunits                                         | 1122 | Cancers Can Evolve Abruptly Due to Genetic Instability                                                                    | 1168 |
| In Plants, Plasmodesmata Perform Many of the Same Functions as Gap Junctions                                   | 1123 | Some Cancers May Harbor a Small Population of Stem Cells                                                                  | 1170 |
| Selectins Mediate Transient Cell–Cell Adhesions in the Bloodstream                                             | 1125 | A Common Set of Hallmarks Typically Characterizes Cancerous Growth                                                        | 1171 |
| Members of the Immunoglobulin Superfamily Mediate Ca <sup>2+</sup> -independent Cell–Cell Adhesion             | 1126 | Cancer Cells Display an Altered Control of Growth and Homeostasis                                                         | 1172 |
| Summary                                                                                                        | 1127 | Human Cancer Cells Escape a Built-in Limit to Cell Proliferation                                                          | 1173 |
| THE EXTRACELLULAR MATRIX OF ANIMALS                                                                            | 1127 | Cancer Cells Have an Abnormal Ability to Bypass Death Signals                                                             | 1174 |
| The Extracellular Matrix Is Made and Oriented by the Cells Within It                                           | 1128 | Cancer Cells Have Altered Sugar Metabolism                                                                                | 1175 |
| Glycosaminoglycan (GAG) Chains Occupy Large Amounts of Space and Form Hydrated Gels                            | 1129 | The Tumor Microenvironment Influences Cancer Development                                                                  | 1175 |
| Hyaluronan Acts as a Space Filler During Tissue Morphogenesis and Repair                                       | 1129 | Cancer Cells Must Survive and Proliferate in a Foreign Environment                                                        | 1176 |
| Proteoglycans Are Composed of GAG Chains Covalently Linked to a Core Protein                                   | 1130 | Summary                                                                                                                   | 1178 |
| Collagens Are the Major Proteins of the Extracellular Matrix                                                   | 1132 | CANCER-CRITICAL GENES: HOW THEY ARE FOUND AND WHAT THEY DO                                                                | 1178 |
| Collagen Chains Undergo a Series of Post-translational Modifications                                           | 1133 | The Identification of Gain-of-Function and Loss-of-Function Cancer Mutations Has Traditionally Required Different Methods | 1179 |
| Secreted Fibril-associated Collagens Help Organize the Fibrils                                                 | 1135 | Retroviruses Led to the Identification of Oncogenes                                                                       | 1180 |
| Elastin Gives Tissues Their Elasticity                                                                         | 1136 | Genes Mutated in Cancer Can Be Made Overactive in Many Ways                                                               | 1181 |
| Cells Govern and Respond to the Mechanical Properties of the Matrix                                            | 1137 | Studies of Rare Hereditary Cancer Syndromes First Identified Tumor Suppressor Genes                                       | 1182 |
| Fibronectin and Other Multidomain Glycoproteins Help Organize the Matrix                                       | 1138 | Both Genetic and Epigenetic Mechanisms Can Inactivate Tumor Suppressor Genes                                              | 1183 |
| Fibronectin Binds to Integrins                                                                                 | 1139 | Systematic Sequencing of Cancer Cell Genomes Has Transformed Our Understanding of the Disease                             | 1184 |
| Tension Exerted by Cells Regulates the Assembly of Fibronectin Fibrils                                         | 1140 | Many Cancers Have an Extraordinarily Disrupted Genome                                                                     | 1185 |
| The Basal Lamina Is a Specialized Form of Extracellular Matrix                                                 | 1141 | Epigenetic and Chromatin Changes Contribute to Most Cancers                                                               | 1185 |
| Laminin and Type IV Collagen Are Major Components of the Basal Lamina                                          | 1141 | Hundreds of Human Genes Contribute to Cancer                                                                              | 1186 |
| Basal Laminae Have Diverse Functions                                                                           | 1143 | Disruptions in a Handful of Key Pathways Are Common to Many Cancers                                                       | 1187 |
| Cells Have to Be Able to Degrade Matrix, as Well as Make It                                                    | 1144 | Mutations in the PI 3-kinase/Akt/mTOR Pathway Drive Cancer Cells to Grow                                                  | 1188 |
| Matrix Proteoglycans and Glycoproteins Regulate the Activities of Secreted Proteins                            | 1145 | Mutations in the p53 Pathway Enable Cancer Cells to Survive and Proliferate Despite Stress and DNA Damage                 | 1189 |
| Summary                                                                                                        | 1146 | Studies Using Mice Help to Define the Functions of Cancer-critical Genes                                                  | 1190 |
| CELL–MATRIX JUNCTIONS                                                                                          | 1147 | Cancers Become More and More Heterogeneous as They Progress                                                               | 1192 |
| Integrins Are Transmembrane Heterodimers That Link the Extracellular Matrix to the Cytoskeleton                | 1147 | Cancers Evolve Slowly Via a Succession of Visible Changes                                                                 | 1192 |
| Integrin Defects Are Responsible for Many Genetic Diseases                                                     | 1148 | A Few Key Genetic Lesions Are Common to a Large Fraction of Colorectal Cancers                                            | 1194 |
| Integrins Can Switch Between an Active and an Inactive Conformation                                            | 1149 | Some Colorectal Cancers Have Defects in DNA Mismatch Repair                                                               | 1195 |
| Integrins Cluster to Form Strong Adhesions                                                                     | 1151 | The Steps of Tumor Progression Can Often Be Correlated with Specific Mutations                                            | 1196 |
| Extracellular Matrix Attachments Act Through Integrins to Control Cell Proliferation and Survival              | 1151 | The Changes in Tumor Cells That Lead to Metastasis Are Still Largely a Mystery                                            | 1197 |
| Integrins Recruit Intracellular Signaling Proteins at Sites of Cell–Matrix Adhesion                            | 1152 | Summary                                                                                                                   | 1197 |
| Cell–Matrix Adhesions Respond to Mechanical Forces                                                             | 1153 | CANCER PREVENTION AND TREATMENT: PRESENT AND FUTURE                                                                       | 1198 |
| Summary                                                                                                        | 1154 | Epidemiology Reveals That Many Cases of Cancer Are Preventable                                                            | 1198 |
| THE PLANT CELL WALL                                                                                            | 1154 | Sensitive Assays Can Detect Those Cancer-causing Agents That Damage DNA                                                   | 1199 |
| The Composition of the Cell Wall Depends on the Cell Type                                                      | 1155 | Fifty Percent of Cancers Could Be Prevented by Changes in Lifestyle                                                       | 1200 |
| The Tensile Strength of the Cell Wall Allows Plant Cells to Develop Turgor Pressure                            | 1155 | Viruses and Other Infections Contribute to a Significant Proportion of Human Cancers                                      | 1201 |
| The Primary Cell Wall Is Built from Cellulose Microfibrils Interwoven with a Network of Pectic Polysaccharides | 1156 | Cancers of the Uterine Cervix Can Be Prevented by Vaccination Against Human Papillomavirus                                | 1202 |
| Oriented Cell Wall Deposition Controls Plant Cell Growth                                                       | 1157 | Infectious Agents Can Cause Cancer in a Variety of Ways                                                                   | 1203 |
| Microtubules Orient Cell Wall Deposition                                                                       | 1158 | The Search for Cancer Cures Is Difficult but Not Hopeless                                                                 | 1204 |
| Summary                                                                                                        | 1159 | Traditional Therapies Exploit the Genetic Instability and Loss of Cell-Cycle Checkpoint Responses in Cancer Cells         | 1204 |
| Problems                                                                                                       | 1160 | New Drugs Can Kill Cancer Cells Selectively by Targeting Specific Mutations                                               | 1204 |
| References                                                                                                     | 1162 | PARP Inhibitors Kill Cancer Cells That Have Defects in <i>Brca1</i> or <i>Brca2</i> Genes                                 | 1205 |
| Chapter 20 Cancer                                                                                              | 1163 |                                                                                                                           |      |
| CANCER AS A MICROEVOLUTIONARY PROCESS                                                                          | 1163 |                                                                                                                           |      |
| Cancer Cells Bypass Normal Proliferation Controls and Colonize Other Tissues                                   | 1164 |                                                                                                                           |      |
| Most Cancers Derive from a Single Abnormal Cell                                                                | 1165 |                                                                                                                           |      |

|                                                                                                                               |             |                                                                                                                                    |             |
|-------------------------------------------------------------------------------------------------------------------------------|-------------|------------------------------------------------------------------------------------------------------------------------------------|-------------|
| Small Molecules Can Be Designed to Inhibit Specific Oncogenic Proteins                                                        | 1207        | A Gene Expression Oscillator Acts as a Clock to Control Vertebrate Segmentation                                                    | 1249        |
| Many Cancers May Be Treatable by Enhancing Immune Responses                                                                   | 1209        | Cell-intrinsic Timing Mechanisms Can Lead to Different Cell Fates                                                                  | 1251        |
| Immunosuppression Is a Major Hurdle for Cancer Immunotherapy                                                                  | 1210        | Cells Rarely Count Cell Divisions to Time Their Development                                                                        | 1252        |
| Cancers Evolve Resistance to Therapies                                                                                        | 1212        | MicroRNAs Can Regulate Developmental Transitions                                                                                   | 1252        |
| We Now Have the Tools to Devise Combination Therapies Tailored to the Individual                                              | 1212        | Cell and Nuclear Size Relationships Schedule the Onset of Zygotic Gene Expression                                                  | 1254        |
| Summary                                                                                                                       | 1213        | Hormonal Signals Coordinate the Timing of Developmental Transitions                                                                | 1255        |
| Problems                                                                                                                      | 1214        | Environmental Cues Determine the Time of Flowering                                                                                 | 1256        |
| References                                                                                                                    | 1216        | Summary                                                                                                                            | 1257        |
| <b>Chapter 21 Development of Multicellular Organisms</b>                                                                      | <b>1217</b> | <b>MORPHOGENESIS</b>                                                                                                               | <b>1257</b> |
| <b>OVERVIEW OF DEVELOPMENT</b>                                                                                                | <b>1218</b> | Imbalance in Physical Forces Acting on Cells Drives Morphogenesis                                                                  | 1258        |
| Conserved Mechanisms Establish the Core Tissues of Animals                                                                    | 1218        | Tension and Adhesion Determine Cell Packing Within Epithelial Sheets                                                               | 1258        |
| The Developmental Potential of Cells Becomes Progressively Restricted                                                         | 1219        | Changing Patterns of Cell Adhesion Molecules Force Cells into New Arrangements                                                     | 1259        |
| Cell Memory Underlies Cell Decision-Making                                                                                    | 1220        | Repulsive Interactions Help Maintain Tissue Boundaries                                                                             | 1259        |
| Several Model Organisms Have Been Crucial for Understanding Development                                                       | 1220        | Groups of Similar Cells Can Perform Dramatic Collective Rearrangements                                                             | 1261        |
| Regulatory DNA Seems Largely Responsible for the Differences Between Animal Species                                           | 1220        | Planar Cell Polarity Orients Cell Behaviors Within an Embryo                                                                       | 1261        |
| Small Numbers of Conserved Cell–Cell Signaling Pathways Coordinate Spatial Patterning                                         | 1221        | An Epithelium Can Bend During Development to Form a Tube                                                                           | 1263        |
| Through Combinatorial Control and Cell Memory, Simple Signals Can Generate Complex Patterns                                   | 1221        | Interactions Between an Epithelium and Mesenchyme Generate Branching Tubular Structures                                            | 1264        |
| Morphogens Are Diffusible Inductive Signals That Exert Graded Effects                                                         | 1222        | The Extracellular Matrix Also Influences Tissue Shape                                                                              | 1265        |
| Lateral Inhibition Can Generate Patterns of Different Cell Types                                                              | 1223        | Cell Migration Is Guided by Environmental Signals                                                                                  | 1266        |
| Asymmetric Cell Division Can Also Generate Diversity                                                                          | 1224        | The Distribution of Migrant Cells Depends on Survival Factors                                                                      | 1267        |
| Initial Patterns Are Established in Small Fields of Cells and Refined by Sequential Induction as the Embryo Grows             | 1225        | Cells Migrate in Groups to Achieve Large-Scale Morphogenetic Movements                                                             | 1268        |
| Developmental Biology Provides Insights into Disease and Tissue Maintenance                                                   | 1225        | Summary                                                                                                                            | 1269        |
| Summary                                                                                                                       | 1226        | <b>GROWTH</b>                                                                                                                      | <b>1269</b> |
| <b>MECHANISMS OF PATTERN FORMATION</b>                                                                                        | <b>1226</b> | The Proliferation, Death, and Size of Cells Determine Organ and Organism Size                                                      | 1270        |
| Different Animals Use Different Mechanisms to Establish Their Primary Axes of Polarization                                    | 1226        | Changes in Cell Size Usually Result from Modified Cell Cycles                                                                      | 1271        |
| Studies in <i>Drosophila</i> Have Revealed Many Genetic Control Mechanisms Underlying Development                             | 1228        | Animals and Organs Can Assess and Regulate Total Cell Mass                                                                         | 1272        |
| Gene Products Deposited in the Egg Organize the Axes of the Early <i>Drosophila</i> Embryo                                    | 1228        | Various Extracellular Signals Stimulate or Inhibit Growth                                                                          | 1273        |
| Three Groups of Genes Control <i>Drosophila</i> Segmentation Along the A-P Axis                                               | 1230        | The Hippo Pathway Relays Mechanical Signals Regulating Growth                                                                      | 1273        |
| A Hierarchy of Gene Regulatory Interactions Subdivides the <i>Drosophila</i> Embryo                                           | 1231        | Hormones Coordinate Growth Throughout the Body                                                                                     | 1274        |
| Egg-Polarity, Gap, and Pair-Rule Genes Create a Transient Pattern That Is Remembered by Segment-Polarity and <i>Hox</i> Genes | 1233        | The Duration of Growth Influences Organism Size                                                                                    | 1275        |
| <i>Hox</i> Genes Permanently Pattern the A-P Axis                                                                             | 1233        | Summary                                                                                                                            | 1275        |
| <i>Hox</i> Proteins Give Each Segment Its Individuality                                                                       | 1234        | Problems                                                                                                                           | 1276        |
| <i>Hox</i> Genes Are Expressed According to Their Order in the <i>Hox</i> Complex                                             | 1234        | References                                                                                                                         | 1278        |
| Trithorax and Polycomb Group Proteins Regulate <i>Hox</i> Expression to Maintain a Permanent Record of Positional Information | 1235        | <b>Chapter 22 Stem Cells in Tissue Homeostasis and Regeneration</b>                                                                | <b>1279</b> |
| The D-V Signaling Genes Create a Gradient of the Transcription Regulator Dorsal                                               | 1236        | <b>STEM CELLS AND TISSUE HOMEOSTASIS</b>                                                                                           | <b>1279</b> |
| A Hierarchy of Inductive Interactions Subdivides the Vertebrate Embryo                                                        | 1238        | Stem Cells Are Defined by Their Ability to Self-renew and Produce Differentiated Cells                                             | 1280        |
| A Competition Between Secreted Signaling Proteins Patterns the Vertebrate Embryonic Axes                                      | 1239        | The Epithelial Lining of the Small Intestine Is Continually Renewed Through Cell Proliferation in Crypts                           | 1281        |
| <i>Hox</i> Genes Control the Vertebrate A-P Axis                                                                              | 1240        | Epidermal Stem Cells Maintain a Self-renewing, Waterproof, Epithelial Barrier on the Body Surface                                  | 1282        |
| Some Transcription Regulators Can Activate a Program That Defines a Cell Type or Creates an Entire Organ                      | 1241        | Cell Lineage Tracing Reveals the Location of Stem Cells and Their Progeny                                                          | 1284        |
| Notch-mediated Lateral Inhibition Refines Cellular Spacing Patterns                                                           | 1242        | Quiescent Stem Cells Are Difficult to Identify by Lineage Tracing                                                                  | 1285        |
| Cell-fate Determinants Can Be Asymmetrically Inherited                                                                        | 1244        | Hematopoietic Stem Cells Can Be Identified by Transplantation                                                                      | 1286        |
| Evolution of Regulatory DNA Explains Many Morphological Differences                                                           | 1245        | Some Tissues Do Not Require Stem Cells for Their Maintenance                                                                       | 1289        |
| Summary                                                                                                                       | 1247        | In Response to Injury, Some Differentiated Cells Can Revert to Progenitor Cells and Some Progenitor Cells Can Revert to Stem Cells | 1289        |
| <b>DEVELOPMENTAL TIMING</b>                                                                                                   | <b>1248</b> | Some Tissues Lack Stem Cells and Are Not Renewable                                                                                 | 1290        |
| Molecular Lifetimes Play a Critical Part in Developmental Timing                                                              | 1249        | Summary                                                                                                                            | 1290        |
|                                                                                                                               |             | <b>CONTROL OF STEM-CELL FATE AND SELF-RENEWAL</b>                                                                                  | <b>1291</b> |
|                                                                                                                               |             | The Stem-Cell Niche Maintains Stem-Cell Self-Renewal                                                                               | 1291        |
|                                                                                                                               |             | The Size of the Niche Can Determine the Number of Stem Cells                                                                       | 1292        |
|                                                                                                                               |             | Asymmetric Stem-Cell Division Can Maintain Stem-Cell Number                                                                        | 1293        |
|                                                                                                                               |             | In Many Symmetric Stem-Cell Divisions, Daughter Cells Choose Their Fates Independently and Stochastically                          | 1294        |
|                                                                                                                               |             | A Decline in Stem-Cell Function Contributes to Tissue Aging                                                                        | 1294        |

|                                                                                                                        |             |                                                                                                       |             |
|------------------------------------------------------------------------------------------------------------------------|-------------|-------------------------------------------------------------------------------------------------------|-------------|
| Summary                                                                                                                | 1296        | THE HUMAN MICROBIOTA                                                                                  | 1347        |
| REGENERATION AND REPAIR                                                                                                | 1296        | The Human Microbiota Is a Complex Ecological System                                                   | 1347        |
| Planarian Flatworms Contain Stem Cells That Can Regenerate a Whole New Body                                            | 1297        | The Microbiota Influences Our Development and Health                                                  | 1348        |
| Some Vertebrates Can Regenerate Entire Limbs and Organs                                                                | 1298        | Summary                                                                                               | 1349        |
| Stem Cells Can Be Used Clinically to Replace Lost Hematopoietic or Skin Cells                                          | 1299        | Problems                                                                                              | 1350        |
| Neural Stem Cells Can Be Manipulated in Culture and Used to Repopulate a Diseased Central Nervous System               | 1299        | References                                                                                            | 1351        |
| Summary                                                                                                                | 1300        |                                                                                                       |             |
| CELL REPROGRAMMING AND PLURIPOTENT STEM CELLS                                                                          | 1300        | <b>Chapter 24 The Innate and Adaptive Immune Systems</b>                                              | <b>1353</b> |
| Nuclei Can Be Reprogrammed by Transplantation into Foreign Cytoplasm                                                   | 1301        | THE INNATE IMMUNE SYSTEM                                                                              | 1354        |
| Reprogramming of a Transplanted Nucleus Involves Drastic Changes in Chromatin                                          | 1301        | Epithelial Surfaces Serve as Barriers to Infection                                                    | 1354        |
| Embryonic Stem (ES) Cells Can Generate Any Part of the Body                                                            | 1302        | Pattern Recognition Receptors (PRRs) Recognize Conserved Features of Pathogens                        | 1354        |
| A Core Set of Transcription Regulators Defines and Maintains the ES-Cell State                                         | 1303        | There Are Multiple Families of PRRs                                                                   | 1355        |
| Fibroblasts Can Be Reprogrammed to Create Induced Pluripotent Stem (iPS) Cells                                         | 1303        | Activated PRRs Trigger an Inflammatory Response at Sites of Infection                                 | 1356        |
| Reprogramming Involves a Massive Upheaval of the Gene Control System                                                   | 1304        | Phagocytic Cells Seek, Engulf, and Destroy Pathogens                                                  | 1358        |
| An Experimental Manipulation of Factors That Modify Chromatin Can Increase Reprogramming Efficiencies                  | 1305        | Complement Activation Targets Pathogens for Phagocytosis or Lysis                                     | 1358        |
| ES and iPS Cells Can Be Guided to Generate Specific Adult Cell Types and Even Organoids                                | 1306        | Virus-infected Cells Take Drastic Measures to Prevent Viral Replication                               | 1360        |
| Cells of One Specialized Type Can Be Forced to Transdifferentiate Directly into Another                                | 1306        | Natural Killer Cells Induce Virus-infected Cells to Kill Themselves                                   | 1361        |
| ES and iPS Cells Are Also Useful for Drug Discovery and Analysis of Disease                                            | 1308        | Dendritic Cells Provide the Link Between the Innate and Adaptive Immune Systems                       | 1362        |
| Summary                                                                                                                | 1309        | Summary                                                                                               | 1362        |
| Problems                                                                                                               | 1310        | OVERVIEW OF THE ADAPTIVE IMMUNE SYSTEM                                                                | 1364        |
| References                                                                                                             | 1312        | B Cells Develop in the Bone Marrow, T Cells in the Thymus                                             | 1365        |
|                                                                                                                        |             | Immunological Memory Depends on Both Clonal Expansion and Lymphocyte Differentiation                  | 1366        |
|                                                                                                                        |             | Most B and T Cells Continually Recirculate Through Peripheral Lymphoid Organs                         | 1368        |
|                                                                                                                        |             | Immunological Self-tolerance Ensures That B and T Cells Do Not Attack Normal Host Cells and Molecules | 1370        |
|                                                                                                                        |             | Summary                                                                                               | 1372        |
| <b>Chapter 23 Pathogens and Infection</b>                                                                              | <b>1313</b> | B CELLS AND IMMUNOGLOBULINS                                                                           | 1372        |
| INTRODUCTION TO PATHOGENS                                                                                              | 1313        | B Cells Make Immunoglobulins (Igs) as Both Cell-Surface Antigen Receptors and Secreted Antibodies     | 1373        |
| Pathogens Can Be Viruses, Bacteria, or Eukaryotes                                                                      | 1314        | Mammals Make Five Classes of Igs                                                                      | 1373        |
| Pathogens Interact with Their Hosts in Different Ways                                                                  | 1314        | Ig Light and Heavy Chains of Antibodies Consist of Constant and Variable Regions                      | 1375        |
| Bacteria Are Diverse and Occupy a Remarkable Variety of Ecological Niches                                              | 1315        | Ig Genes Are Assembled from Separate Gene Segments During B Cell Development                          | 1377        |
| Bacterial Pathogens Carry Specialized Virulence Genes                                                                  | 1317        | Antigen-driven Somatic Hypermutation Fine-Tunes Antibody Responses                                    | 1379        |
| Bacterial Virulence Genes Encode Toxins and Secretion Systems That Deliver Effector Proteins to Host Cells             | 1319        | B Cells Can Switch the Class of Ig They Make                                                          | 1379        |
| Fungal and Protozoan Parasites Have Complex Life Cycles Involving Multiple Forms                                       | 1321        | Summary                                                                                               | 1381        |
| All Aspects of Viral Propagation Depend on Host-Cell Machinery                                                         | 1322        | T CELLS AND MHC PROTEINS                                                                              | 1382        |
| Summary                                                                                                                | 1325        | T Cell Receptors (TCRs) Are Ig-like Heterodimers                                                      | 1382        |
| CELL BIOLOGY OF PATHOGEN INFECTION                                                                                     | 1325        | Activated Dendritic Cells Activate Naïve T Cells                                                      | 1383        |
| Pathogens Breach Epithelial Barriers to Infect the Host                                                                | 1326        | T Cells Recognize Foreign Peptides Bound to MHC Proteins                                              | 1384        |
| Pathogens That Colonize an Epithelium Must Overcome Its Protective Mechanisms                                          | 1326        | MHC Proteins Are the Most Polymorphic Human Proteins Known                                            | 1388        |
| Extracellular Pathogens Use Toxins and Contact-dependent Secretion Systems to Disturb Host Cells Without Entering Them | 1328        | CD4 and CD8 Co-receptors on T Cells Bind to Invariant Parts of MHC Proteins                           | 1389        |
| Intracellular Pathogens Have Mechanisms for Both Entering and Leaving Host Cells                                       | 1329        | Developing Thymocytes Undergo Positive and Negative Selection                                         | 1389        |
| Viruses Bind to Virus Receptors at the Host-Cell Surface                                                               | 1329        | Cytotoxic T Cells Induce Infected Target Cells to Undergo Apoptosis                                   | 1391        |
| Viruses Enter Host Cells by Membrane Fusion, Pore Formation, or Membrane Disruption                                    | 1330        | Effector Helper T Cells Help Activate Other Cells of the Innate and Adaptive Immune Systems           | 1392        |
| Bacteria Enter Host Cells by Phagocytosis                                                                              | 1331        | Naïve Helper T Cells Can Differentiate into Different Types of Effector T Cells                       | 1393        |
| Intracellular Eukaryotic Parasites Actively Invade Host Cells                                                          | 1333        | Both T and B Cells Require Multiple Extracellular Signals for Activation                              | 1394        |
| Some Intracellular Pathogens Escape from the Phagosome into the Cytosol                                                | 1334        | Many Cell-Surface Proteins Belong to the Ig Superfamily                                               | 1396        |
| Many Pathogens Alter Membrane Traffic in the Host Cell to Survive and Replicate                                        | 1335        | Vaccination Against Pathogens Has Been Immunology's Greatest Contribution to Human Health             | 1396        |
| Bacteria and Viruses Use the Host-Cell Cytoskeleton for Intracellular Movement                                         | 1338        | Summary                                                                                               | 1400        |
| Many Microbes Manipulate Autophagy                                                                                     | 1340        | Problems                                                                                              | 1402        |
| Viruses Can Take Over the Metabolism of the Host Cell                                                                  | 1340        | References                                                                                            | 1404        |
| Pathogens Can Evolve Rapidly by Antigenic Variation                                                                    | 1341        |                                                                                                       |             |
| Error-prone Replication Dominates Viral Evolution                                                                      | 1343        |                                                                                                       |             |
| Drug-resistant Pathogens Are a Growing Problem                                                                         | 1344        |                                                                                                       |             |
| Summary                                                                                                                | 1346        |                                                                                                       |             |
|                                                                                                                        |             | <b>Glossary</b>                                                                                       | <b>G:1</b>  |
|                                                                                                                        |             | <b>Index</b>                                                                                          | <b>I:1</b>  |